POLYHEDRON

e International Journal for Inorganic and Organometallic Chemistry

incorporating *Journal of Inorganic and Nuclear Chemistry*
and *Inorganic and Nuclear Chemistry Letters*

Editor-in-Chief
Sir Geoffrey Wilkinson FRS

Executive Editors
D. C. Bradley FRS
M. H. Chisholm

UNIVERSITY OF HAWAII
LIBRARIES
AUG 1 8 1983

Subject Index
Volume 1, 1982

Compiled by Peter Thornton

LIBRARY USE
ONLY

ER
D
64

Pergamon Press

Oxford ● New York ● Paris ● Frankfurt ● Toronto ● Sydney

POLYHEDRON

The International Journal for Inorganic and Organometallic Chemistry
(incorporating *Journal of Inorganic and Nuclear Chemistry* and *Inorganic and Nuclear Chemistry Letters*)

Editor-in-Chief:
Sir Geoffrey Wilkinson, FRS, *Imperial College of Science and Technology*

Executive Editors:
Rest of World
D C Bradley, FRS, *Queen Mary College, University of London, Mile End Road, London E1 4NS, UK*

U.S.A. and Canada
M H Chisholm, *Department of Chemistry, Indiana University, Bloomington, IN 47405, U.S.A.*

Editorial Board
E W Abel, *University of Exeter*
J E Baldwin, FRS, *University of Oxford*
R D Gillard, *University College, Cardiff*
M L H Green, *University of Oxford*
M B Hursthouse, *Queen Mary College, University of London*
N N Greenwood, *University of Leeds*
Sir Jack Lewis, FRS, *University of Cambridge*
S F Mason, FRS, *King's College, University of London*
W McFarlane, *City of London Polytechnic*
B L Shaw, FRS, *University of Leeds*
F G A Stone, FRS, *University of Bristol*
A G Sykes, *University of Newcastle upon Tyne*
J M Thomas, FRS, *University of Cambridge*
K Wade, *University of Durham*

Polyhedron Reports and Polyhedron Symposia-in-Print
Editors:
M H Chisholm, *Indiana University*
B F G Johnson, *University of Cambridge*

Publishing Office: European Publishing Corporation Ltd., 84–86 Malahide Road, Coolock, Dublin 5, Republic of Ireland (phone Dublin 475377; telex 25670)

Subscription and Advertising Offices: Headington Hill Hall, Oxford OX3 0BW (phone Oxford 64881; telex 83177). Maxwell House, Fairview Park, Elmsford, NY 10523, USA (phone 914 592 7700; telex 137328)

Twelve issues per year

Annual Subscription Rates (1983)
For libraries, university departments, government laboratories, industrial and other multiple-reader institutions US $500.00 per annum (including postage and insurance). Two year rate (1983–84) US $950.00.

Specially Reduced Rates to Individuals
In the interests of maximising the dissemination of the research results published in this important international journal we have established a two-tier price structure. Any individual whose institution takes out a library subscription may purchase a second or additional subscription for personal use at the much reduced rate of US $55.00.

Microform Subscriptions and Back Issues
Back issues of all previously published volumes are availabe in the regular editions and on microfilm and microfiche. Current subscriptions are available on microfiche simultaneously with the paper edition and on microfilm on completion of the annual index at the end of the subscription year.

Copyright © 1983 Pergamon Press Ltd.
It is a condition of publication that manuscripts submitted to this journal have not been published and will not be simultaneously submitted or published elsewhere. By submitting a manuscript, the authors agree that the copyright for their article is transferred to the publisher if and when the article is accepted for publication. However, assignment of copyright is not required from authors who work for organisations which do not permit such assignment. The copyright covers the exclusive rights to reproduce and distribute the article, including reprints, photographic reproductions, microform or any other reproductions of similar nature and translations. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, electrostatic, magnetic tape, mechanical, photocopying, recording or otherwise, without permission in writing from the copyright holder.

Photocopying Information for Users in the U.S.A.
The Item-Fee Code for this publication indicates that authorization to photocopy items for internal or personal use is granted by the copyright holder for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service provided the stated fee for copying beyond that permitted by Section 107 or 108 of the United States Copyright Law is paid. The appropriate remittance of $3.00 per copy per article is paid directly to the Copyright Clearance Center Inc., 21 Congress Street, Salem, MA 01970. The copyright owner's consent does not extend to copying for general distribution, for promotion, for creating new works, or for resale. Specific written permission must be obtained from the publisher for such copying. In case of doubt please contact your nearest Pergamon office.

The Item-Fee Code for this publication is: 0277–5387/83 $3.00 + .00

POLYHEDRON

SUBJECT INDEX

Compiled by Peter Thornton

A

Absolute configuration, molybdenum(II) carbonyl cyclopentadienyl pyrrole Schiff base complex, 617
Acetate complexes
  chromium(II), 135, 209
  manganese(II), 327
  molybdenum(II), 209
  osmium(VI), 840
  rhenium(V), 31
  rhodium(II), 637
Acetic acid, solvates, 633
Acetone, formation from propan-2-ol, 839
Acetonitrile, solvent extraction of chromium(III) complexes, 383
Acetonitrile complexes
  copper(I) and (II), 467
  molybdenum(0), 130
  molybdenum(IV), 209
  rhodium(III), 577
  silver(I), 145
  tungsten(IV), 209
Acetonitrile, NMR (^{13}C) spectrum, effects of added electrolytes, 145
Acetoxytrihydroborate complex, copper(I), 299
Acetyl complexes
  iron(0), 842
  osmium(II), 83
Acetylacetonate, see 2,4-pentanedionate
2-Acetylthiophene-2′-pyridylhydrazone complex, palladium(II), 179
3-Acetylthiophene-2′-pyridylhydrazone complexes, palladium(II), 179
Actinides, separation from lanthanides, 277
Acyl complexes, iron(0), 842
Adenosine, reaction with chromium(VI), 497
5′-Adenosinemonophosphate, reaction with chromium(VI), 497
5′-Adenosinetriphosphate, reaction with chromium(VI), 497
Alanine complexes
  uranium(VI), 537
  zinc(II), 153
Aldehydes
  branched chain, formation in hydroformylations, 589
  formation, catalysis by rhodium(I) complex, 587
Aliquat, see trioctylmethylammonium chloride
Alkenes
  cyclodimerisation catalysed by metallacycles, 760, 765, 771
  formation of metallacycles, 750
  hydroformylation, kinetic study, 591, 593
  metathesis, 770
Alkoxide complex, iron(0), 842
Alkylammonium salts, synergic effects in solvent extraction, 457
Alkylidyne complexes
  iron(0), 842
  rhenium(I), 803
Alkynes
  formation of metallacycles, 750
  trimerisation catalysed by metallacycles, 756
Allene, dimerisation, 769
Allyl complex
  formation from metallacycle, 747
  rhodium(I), 574
Allyl alcohol, 594
Allylbenzene, isomerisation, 580
Allyl-*t*-butylether, 594
Allyl(2,6-dimethylphenyl)ether, isomerisation, 580
Alumina, hydroformylation catalyst support, 597
Aluminium 3+, reaction with fluoropolytungstate, 263
Aluminium(III)
  bis(pyrogallol dianion)-, ammonium salt monohydrate, 313
  catechol dianion complexes, 313

  chlorodimethyl-, reaction with metallacycles, 762
  3,4-dihydroxybenzoate dianion complexes, 313
  dihydroxy(pyrogallol dianion)-, ammonium salt, 313
  diphenolate complexes, synthesis and formation constants, 313
  heteropolytungstates, 263
  3,3′,4,4′,5,5′-hexahydroxydiphenylsulphone dianion complexes, 313
  methylthymol blue complexes, electronic spectra, potentiometric titrations and formation constants, 405
  pyrogallol dianion complexes, 313
  3,3′,4,4′-tetrahydroxydiphenylsulphone dianion complexes, 313
  tetrachloro-, titanium(II) complex, 393
  tris(pyrogallol dianion)-, ammonium salt tetrahydrate, 313
Aluminofluorotungstate$\{HAlF_3W_{11}O_{36}(H_2O)\}^{5-}$, tetramethylammonium salt hydrate, 263
Aluminofluorotungstate$\{H_2AlF_2W_{11}O_{37}(H_2O)\}^{5-}$, tetramethylammonium salt hydrate, 263
Aluminofluorotungstates, synthesis, NMR (^{19}F) spectra and polarography, 263
Americium(III), complexes with 2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate, 277
Amide complexes,
  magnesium, 421
  rhenium(I), (III) and (V), 31
Amidotriphosphate, formation from ammonia and trimetaphosphate, 425
Aminoacid complexes
  cadmium(II), 153
  manganese(II), 113
  uranium(VI), 537
  zinc(II), 153
2-Aminobenzothiazole complexes, antimony(III) and bismuth(III), 655
4-Amino-3-nitrobenzoic acid, 487
Aminopolycarboxylate complexes, rhodium(III), 659
N-Aminorhodanine complexes, copper(I) and silver(I), 633
Aminothiazoline complex, zinc(II), 153
Aminotrihydroxymethane complex, copper(II), 361
Amino(triphenyl)phosphonium chloride, 535
Ammonia, reaction with
  copper and hydrogen peroxide, 437
  trimetaphosphate, 425
Ammonia complexes
  cobalt(III), 365
  platinum(II), 57, 201
  uranium(VI), 483
Ammonium
  fluoride, 553
  β-octamolybdate pentahydrate, X-ray crystal structure, 17
Anion exchange, polyphosphates, 69
Antimony(III)
  2-aminobenzothiazoletriiodo-, 655
  bromide, 655
  chloride, 655
  cyclophosphamidetrichloro-, 637
  di(2-aminobenzothiazole)trichloro-, 655
  halide thiazole complexes, synthesis, IR spectra and electric conductance, 655
  iodide, 655
  sesqui(2-aminobenzothiazole)tribromo-, 655
Apatite, uranium(VI) in, 343
Apotransferrin complex, copper(II), 361
Aquo, *see* water
Arsenic(V) in heteropolytungstates, 283
Arsenostannotungstate$(HAsSnW_{11}O_{40})^{4-}$, potassium salt hydrate, 283
L-Aspartate complex, terbium(III), 693
Atomisation enthalpies, boron compounds, 701
Azide, reaction with platinum(II) complex, 201
Azide complexes, magnesium, 421

B

Base hydrolysis, cobalt(III) complex, 203
Benzene, solvent extraction of zinc(II) complex, 387
η^6-Benzene complex, titanium(II), 393
Benzoate complex, tungsten(IV), 629

N-Benzothiazolyl-N′-ethylthiourea complexes, 557
N-Benzothiazolyl-N′-methylthiourea complexes, 557
Benzyl alcohol, solvent extraction of chromium(III) complexes, 383
Beryllium
  hexanitratooxotetra-, 820
  oxo nitrate, mass spectrum, 820
Biacetyldihydrazone complexes, manganese(II) and zinc(II), 831
Bicarbonate, reaction with copper(II) and apotransferrin, 361
Binuclear complexes
  aluminium(III), 405
  cadmium(II), 157
  chromium(II), 135, 209
  cobalt(III), 21
  copper(II), 143, 679
  mercury(II), 157
  molybdenum(0), 53
  molybdenum(II), 209
  molybdenum(V), 109, 163
  nickel(II), 409
  osmium(II,III), 83
  rhodium(II), 637
  tungsten(I), 803
  uranium(III), 417
  zinc(II), 157
2,2′-Bipyridyl complexes
  chromium(0), 209
  cobalt(II), 247, 711
  copper(I), 467
  copper(II), 247, 467
  hafnium(0), 209
  manganese(II), 327
  mercury(II), 819
  molybdenum(0) and (IV), 209
  nickel(II), 247, 711
  niobium(0) and rhenium(0), 209
  solvent extraction, 387
  tantalum(0), tungsten(0) and (IV) and vanadium(0) 209
  zinc(II), 387
  zirconium(0), 209
4,4′-Bipyridyl complexes, manganese(II), 327
2,2′-Bipyridyl dioxide complexes, manganese(II), 327
4,4′-Bipyridyl dioxide complexes, manganese(II), 327
2,2′-Bipyrimidine complexes,
  binuclear, 53
  chromium(0), molybdenum(0) and tungsten(0), 53
2,2′-Bipyrimidine anion complexes
  molybdenum(0), 53
3,3′-Bis[(bis{carboxymethyl}amino)methyl]thymolsulphonephthalein, see methylthymol blue
Bis(3,4-diphenylcyclopentadienyl)1,2-diphenylethane complex, rhenium(I), 441
1,2- Bisdiphenylphosphinoethane complexes
  molybdenum(0), 130
  platinum(0), 105
  rhodium(I), 597
Bis(diphenylphosphino)methane complexes, rhodium(I), 809
Bis(diphenylphosphinomethyl)phenylphosphine complex, molybdenum(0), 130
6,6′′-Bismethylhydrazino-4,4′′-diphenyl-2,2′:6′,2′′-terpyridyl, synthesis and mass spectrum, 303
6,6′′-Bismethylhydrazino-4′-phenyl-2,2′:6′,2′′-terpyridyl, synthesis, mass spectrum and complexes with 3d-elements, 303
6,6′′-Bismethylhydrazino-4′-phenyl-2,2′:6′,2′′-terpyridyl - glyoxal macrocycle, cobalt(II) complex, 303
Bismuth(III)
  2-aminobenzothiazoletribromo-, 655
  2-aminobenzothiazoletrichloro-, 655
  2-aminobenzothiazoletriiodo-, 655
  bromide, 655
  chloride, 655
  halide thiazole complexes, synthesis, electric conductance and IR spectra, 655
  iodide, 655

Bis(pentafluorophenyl)acetylene, formation of metallacycles, 757
Bistrimethylsilylamide, europium(III), gadolinium(III), indium(III), lanthanum(III) and praseodymium(III) complexes, 307
Blood plasma, copper(II) histamine complexes, 371
Bond enthalpies, boron compounds, 701
Bond lengths (not determined by diffraction methods), germanium and tin organo halides, 397
Book review
  NMR (^{13}C) spectra of organometallic compounds, 139
  organotin compounds, 845
Borane
  di(pyrrolyl-1)(pyridine)-, 673
  pyridine(pyrrolyl-1)-, 673
Boranes, molecular orbital calculation, 521
Borate
  dihydrodi(pyrrolyl-1)-, potassium salt, 673
  hydro(pyrrolyl)-, hydrolysis, kinetic study and isotope exchange, reaction with pyridine, 673
  hydrotri(pyrrolyl-1)-, sodium salt and its dioxane solvate, 673
  pyrrolyl-1-trihydro-, sodium salt dioxane solvate, 673
Borazine, 701
Borogermanotungstate$(HBGeW_{11}O_{40})^{6-}$, potassium salt hydrate, 283
Borohydride, *see* hydroborate
Boric acid, 701
Boron
  compounds, bond enthalpies, atomisation enthalpies and enthalpies of formation, 701
Boron (cont)
  heteropolytungstates, 283
  monobromide, 701
  monochloride, 701
  monofluoride, 701
  oxides, 701
  subhalides, synthesis, mass spectra and reactions, 134, 319
  tribromide, 701
  trichloride, 701
  trifluoride, 701
Borostannotungstate$(HBSnW_{11}O_{40})^{6-}$, potassium salt hydrate, 283
Bromide, reaction with platinum(II) complex, 201
Bromides and bromo complexes
  antimony(III) and bismuth(III), 655
  boron cage compounds, 134
  cadmium(II), 137, 781
  cobalt(II), 377
  copper(I), 633
  copper(II), 377, 543
  germanium, 397
  lanthanum(III), 49
  lead(II), 169
  mercury(II), 137, 543, 667, 781
  neodymium(III), 49
  nickel(II), 377
  palladium(II), 179, 289
  phosphorus(V), 123
  platinum(II), 289, 689
  rhodium(III), 577
  sulphur(VI), 723
  tin(IV), 397
Bromoborane, 701
Bromo(di-*n*-butyl)borane, 701
7-Bromo-5,6-dicarba-*nido*-decaborane(12), 822
Bromo(dimethyl)borane, 701
Bromo(diphenyl)borane, 701
8-Bromododecahydro-*nido*-undecaborate(2) complexes, cobalt(III), 511
1-(2-Bromopyridylcarbonylmethyl)pyridinium iodide, synthesis and NMR (^1H) spectrum, 303
1-(2-Bromopyridyl)-3-phenyl-2-propenone, synthesis and IR and mass spectra, 303
1-Bromo-2-(4-sulphophenyl)ethane, sodium salt, 61
N-Bromotetrafluoromethylamine, 129
Buta-1,3-diene
  formation of metallacycles, 768
  hydrosilylation, 600

Buta-1,3-diene complex, rhenium(V), 37
But-1-ene
  hydroformylation, 591
  isomerisation, 579
But-1-en-3-ol, hydroformylation, 599
But-2-ene, hydroformylation, 599
t-Butoxide complex, tungsten(VI), 561
n-Butyl compounds
  phosphorus(V), 123
t-Butyl compounds
  mercury(II), 413
  tin(IV), 1
Butylamine, carbonylation, 601
t-Butylamine complexes, tungsten(VI), 561
t-Butylaminomethylidyne complexes, rhenium(I), 803
n-Butyldibromoborane, 701
n-Butyldichloroborane , 701
n-Butyldiphenylphosphine complexes, cobalt(-I) and iron(-II), 45
N,*N*′-Butylenebis(2,4-pentanedioneimine) complexes, copper(II), nickel(II), palladium(II), 785
N,*N*′-Butylene[(2,4-pentanedioneimine)(5,5,5-trifluoro-2,4-pentanedioneimine)] complexes, copper(II), nickel(II), palladium(II), 785
N,*N*′-Butylenebis(5,5,5-trifluoro-2,4-pentanedioneimine) complexes, copper(II), nickel(II), palladium(II), 785
t-Butylisocyanide complexes, chromium(0), platinum(II), rhenium(I), tungsten(0) and (I), 803
(4-*t*-Butyl)methylenecyclohexane, hydrogenation, 583
n-Butyl(*i*-propyl)-*n*-propylborane, 701
But-2-yne, 593, 599
Butyraldehyde, 585, 591
Butyrate complex, rhodium(II), 637

C

Cadmium(II)
  arsine carboxylate complexes, synthesis, IR spectra and thermogravimetric analysis, 667
  bis(*N*-benzothiazolyl-*N*′-ethylthiourea)-, 557
  bis(*N*-benzothiazolyl-*N*′-methylthiourea)-, 557
  bis(2-dicyclohexylarsinobenzoato)-, 667
  bis(2-diethylarsinobenzoato)-, 667
  bis-μ-(1-methyl-4-mercaptopiperidine)tetrabromodi-, monohydrate, 157
  bis-μ-(1-methyl-4-mercaptopiperidine)tetraiododi- , monohydrate, 157
  bis(tri-4-tolylphosphine selenide)tetraiododi-, 781
  bis(tri-4-tolylphosphine sulphide)tetraiododi-, 781
  bromide, 781
  bromotris(diphenylthiophosphinyl)methanido-, 137
  chlorotris(diphenylthiophosphinyl)methanido-, 137
  cystinate complexes, formation constants, 153
  diacetatotriaquo-, 557
  dichloro(1-methyl-4-mercaptopiperidine)-, 157
  4′,4‴-diphenyl-2,2′:6′,2″:6″,2‴:6‴,2⁗-quinquepyridine-, hexafluorophosphate dihydrate, 311
  ethylenediaminediacetate complex, 153
  ethylenediaminetetraacetate complex, 153
  halide compounds, preparation, 137
  halide phosphine selenide and phosphine sulphide complexes, synthesis, electric conductance and IR spectra, 781
  halide thiolate complexes, synthesis, IR spectra, electric conductance and X-ray diffraction, 157
  hexakis(tri-4-tolylphosphine selenide)tetrabromodi-, 781
  iodide, 781
  iodo(trifluoromethyl)-, 826
  iodotris(diphenylthiophosphinyl)methanido-, 137
  penicillaminate complex, 153
Caesium
  bromoiodoiodide, 215
  cobalt(III) carbaborane salts, 511
  dibromoiodide, 215

  dichloroiodide, 215
  fluoride, 553
  perbromate, X-ray crystal structure, erratum, 567
  tribromide, 215
  trihalides, thermogravimetric analysis and dissociation enthalpies, 215
  triiodide, 215
Camphorquinone dioxime dianion complex, nickel(II), 503
Camphorquinone dioxime, isomerism, 503
Camphorquinone dioxime monoanion complexes, copper(II) and nickel(II), 503
Carbaboranes
  optical activity and NMR (^{11}B) spectra, 824
  synthesis and NMR (^{11}B) spectra, 822
Carbaborane complexes
  cobalt(III), 511
  iron(II) and ruthenium(II), 77
closo-1-Carba-2-(η^5-cyclopentadienyl)-2-cobaltaoctaborane(9), 833
closo-1-Carba-6-(η^5-cyclopentadienyl)-6-nickelaoctaborane(9), 833
closo-1-Carba-10-(η^5-cyclopentadienyl)-10-nickelaoctaborane(9), 833
Carbon-carbon bonds, insertion of metals, 739
Carbon-hydrogen bond activation
  rhodium(I) complexes, 501
  tungsten compounds, 629
Carbon dioxide complex, rhenium(I), 441
Carbon monoxide
  promoter of selective hydroformylation, 591
  reaction with metallacycles, 749, 756, 763
Carbon tetrachloride, in solvent extraction of aqueous osmium(VI) and (VIII) solutions, 175
Carbonyl complexes
  chromium(0), 53, 819
  cobalt(-I), 45
  iron(-II), 45
  iron(0), 521, 842
  iron(I), 1
  molybdenum(0), 53, 130, 819
  molybdenum(II), 233, 617
  osmium cluster hydrides, 105
  platinum(II) and (IV), 689
  rhenium(I), 37, 441
  rhodium(I), 501, 569, 839
  ruthenium(II), 97
  tungsten(0), 53, 819
  vanadium(0), 209
Carbonylation, catalysis by rhodium(I) complex, 601
Carboxylate complexes
  chromium(II), 135
  cobalt(III), 149
  copper(II), 679
  osmium(VI), 840
  rhenium(IV), 37
  rhenium(V), 31, 37
  rhodium(II), 637
  tellurium(IV), 13
  tungsten(IV), 629
N-(2′-Carboxyphenyl)iminodiacetate complexes, thorium(IV) and uranium(VI), 827
2-Carboxypyridine complex, platinum(II), 57
Carcinogenicity, chromium(VI) compounds, 497
Catalysis
  alkene cyclodimerisation by metallacycles, 760, 765
  alkyne trimerisation by metallacycles, 756
  dehydration of alcohols by zirconium(IV) sulphophosphonates, 61
  esterification of acids by zirconium(IV) sulphophosphonates, 61
  hydrogen peroxide decomposition by osmium(VIII) oxide, 175
  rhodium(I) complex in hydrogenation, hydroformylation, 569
Catechol dianion complexes, aluminium(III), 313
Cation exchange
  separation of actinides and lanthanides, 277
  zirconium(IV) sulphophosphonates, 61
Cavity formation energy, in solvent extraction of chromium(III) β-diketonate complexes, 383

- Cellulose triacetate, use in chromatographic resolution of organotin compound, 1
- Cerium(III)
  - complexes with 2,2′-diaminodiethylether,*N*,*N*,*N′*,*N′*-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N′*,*N′*-tetraacetate, 277
  - difluorohydroxo-, 239
  - dihydroxofluoro-, 239
  - fluorohydroxo-, 239
- Chemical shift anisotropy, relaxation of ^{195}Pt in NMR (^{1}H) spectra of platinum(II) complexes, 57
- Chiral aldehydes, synthesis using rhodium(I) complex, 580
- Chiral phosphines, rhodium(I) complexes, 598
- Chloramines, 535
- Chlorides and chloride complexes
  - aluminium(III), 393
  - antimony(III), 637, 655
  - binuclear, 409
  - bismuth(III), 655
  - boron cage compounds, 134
  - cadmium(II), 137, 157
  - chromium(III), 209, 471, 475
  - cobalt(II), 21, 377, 623
  - cobalt(III), 525
  - copper(I), 633
  - copper(II), 365, 377, 543
  - effect on copper(II)-iodide reaction, 611
  - gallium(III), 623
  - germanium(IV), 397
  - hafnium(IV), 209
  - indium(III), 355, 457, 549, 623
  - iridium(III), 525
  - iron(III), 623
  - lanthanum(III), 49
  - manganese(II), 327, 831
  - mercury(II), 137, 157, 243, 543, 667, 781
  - molybdenum(0), 130
  - molybdenum(IV), 209
  - neodymium(III), 49
  - nickel(II), 377, 409
  - niobium(V), 209
  - osmium(II), (III), (IV) and (II,III), 83
  - palladium(II), 179, 289, 543
  - phosphorus(V), 123
  - platinum(II), 57, 89, 201, 289, 525, 689
  - platinum(IV), 57, 525, 689
  - rhenium(I), 441, 803
  - rhenium(III), 441
- Chlorides and chloride complexes (cont)
  - rhenium(IV), 209
  - rhenium(V), 31
  - rhodium(I), 501, 575, 809
  - rhodium(III), 525, 543
  - ruthenium(II), 525
  - ruthenium(III), 525, 543
  - silver(I), 97
  - solvent extraction with tetraalkylammonium salts, dielectric constants, electric conductance, IR spectra and viscosities, 623
  - tantalum(V), 209
  - tin(IV), 209, 397
  - titanium(II), 393
  - titanium(IV), 209
  - tungsten(IV), 209, 629
  - tungsten(VI), 561
  - uranium(IV), 417
  - vanadium(III), 209
  - zinc(II), 157, 543, 623, 831
  - zirconium(IV), 209
- 4-Chloroacetophenone-2′-pyridylhydrazone complexes, palladium(II), 179
- Chloroborane, 701
- 3-Chlorobutene, isomerisation, 581
- Chloro(di-*n*-butyl)borane, 701

4-Chloro-5,6-dicarba-*nido*-decaborane(12), 822
8-Chloro-5,6-dicarba-*nido*-decaborane(12), 822
10-Chloro-5,6-dicarba-*nido*-decaborane(12), 822
Chlorodimethylborane, 701
Chlorodimethylsilane, 601
Chloro(diphenyl)borane, 701
Chloro(diphenyl)phosphine, 535
8-Chlorododecahydrodicarba-*nido*-undecaborate(2) complexes, cobalt(III), 511
9-Chlorododecahydrodicarba-*nido*-undecaborate(2) complexes, cobalt(III), 511
1-Chlorotetrafluoroethyl compound, sulphur(VI), 723
Chromatography, in resolution of enantiomers of organotin compound, 1
Chromium(0)
  bipyrimidine bridged binuclear complex, 53
  μ-2,2′-bipyrimidine(octacarbonyl)molybdenum-, 53
  carbonyl phosphine complexes, synthesis and NMR (^{31}P) spectra, 819
  hexa-*t*-butylisocyanide-, 803
  isocyanide complexes, synthesis, IR and NMR (^{1}H, ^{13}C) spectra, 803
  tris(2,2′-bipyridyl)-, 209
  tris(1,10-phenanthroline)-, 209
Chromium(II)
  binuclear acetate thiocyanate complex, synthesis, electronic and IR spectra and magnetic properties, 135
  tetra-μ-acetatodi-, 209, 803
  tetra-μ-acetatodiisothiocyanatodi-, tetraethylammonium salt, 135
Chromium(III)
  aquo-*trans*-bis(methylnicotinate)-*mer*-trichloro-, 471
  bis[1,2-dimethylnonahydrodicarba-*nido*-undecaborate(2)]- caesium salt monohydrate, 663
  6,6′′-bismethylhydrazino-4′-phenyl-2,2′:6′,2′′-terpyridyl-, chloride hydrochloride, 303
  bis[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
  chloride, 209
  chloride, IR and Raman spectra and factor group analysis, 475
  chloride methylnicotinate water complex, synthesis and X-ray crystal structure, 471
  β-diketonate complexes, isomers, solvent extraction, 383
  hexathiocyanato-, electronic spectrum and solvolysis, 191
  thiocyanate complex, electronic spectra solvent dependence and solvolysis, 191
  tris(1,1,1-trifluoro-2,4-pentanedionato)-, isomers, 383
Chromium(V), formation in reaction of chromium(VI) with ribonucleotides, ESR spectra, 497
Chromium(VI)
  reaction with ribonucleotides and carcinogenicity, 497
  tetroxo- sodium salt, 497
Circular dichroism, rhodium(III) aminocarboxylate diamine complex, 659
Circularly polarised fluorescence spectra, lanthanide complexes, 195, 693
Cisplatin, *see* platinum(II), *cis*-diamminodichloro-
Cluster compounds
  cobalt boron cyclopentadienyl hydride, 133
  dodecahedral, bonding scheme and violation of Wade's rules, 133
  iron(0), 842
  nickel boron cyclopentadienyl hydride, 133
  osmium gold carbonyl hydrides, 105
Cobalamins, NMR (^{1}H) spectra, spin-lattice relaxation, conformation changes and TLC, 321
Cobalt
  carbaborane complex, synthesis and NMR (^{1}H, ^{11}B) spectra, 833
  η^{5}-cyclopentadienyl borane complex, bonding, 133
Cobalt(-I)
  bis(tetracarbonyl-)zinc(II), reactions with unsaturated compounds, NMR (^{1}H), ^{13}C) spectra, 563
  compound with cobalt-lead bond, 45
Cobalt(I)
  η^{3}-allyltricarbonyl-, 563
  cyclopentadienyldicarbonyl-, 663, 743
  dicarbonyl(pentamethylcyclopentadienyl)-, 663
  dicarbonyl[undecahydrodicarba-*nido*-undecaborate(2)]-, 663

  - 4′,4‴-diphenyl-2,2′:6′,2″:6″,2‴:6‴,2⁗-quinquepyridyl complex, 311
  - hydridotetracarbonyl-, 585
- Cobalt (2+), reaction with fluoropolytungstate, 263
- Cobalt(II)
  - bis(*N*-benzothiazolyl-*N*′-ethylthiourea)-, 557
  - bis(*N*-benzothiazolyl-*N*′-methylthiourea)-, 557
- Cobalt(II) (cont)
  - bis(2,2′-bipyridyl)nitroacetato-, dihydrate, 247
  - bisdimethylsulphoxidedi[1-naphthylmercury(II)thiocyanato]-dithiocyanato-, 711
  - bisdimethylsulphoxidedithiocyanatodi[4-tolylmercury(II)-thiocyanato]-, 711
  - 6,6″-bismethylhydrazino-4′-phenyl-2,2′:6′,2″-terpyridyl-, tetrafluoroborate methanol solvate, 303
  - (6,6″-bismethylhydrazino-4′-phenyl-2,2′:6′,2″-terpyridyl-glyoxal macrocycle) hexafluorophosphate dihydrate, 303
  - bispentamethylcyclopentadienyl-, 663
  - bis(1-phenyl-3-methyl-4-benzoylpyrazol-5-onato)-, 461
  - bispyridinedi[1-naphthylmercury(II)thiocyanato]dithiocyanato-, 711
  - bispyridinedithiocyanato-, 711
  - bispyridinedithiocyanatodi[4-tolylmercury(II)thiocyanato]-, 711
  - bis[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
  - chlorobis(di-2-pyridylketone)pyridine-, chloride tetrahydrate, 21
  - cobalamins and cobinamides, 321
  - diacetatotetraaquo-, 557
  - diaquodichloro-, 543
  - 2,6-dibenzothiazol-2-ylpyridinediiodo-, 543
  - dibromo(Me_4-[16]-tetene)-, 377
  - dichloro(Me_4-[16]-tetene)-, 377
  - dicyclopentadienyl-, 663
  - dihistidine-, 791
  - dihistidine(pyridoxamine)-, 791
  - β-diketonate complexes, formation constants, solvent extraction and synergic effects, 461
  - di[1-naphthylmercury(II)thiocyanato]dithiocyanato-, 711
  - dinitrato(Me_4-[16]-tetene)-, 377
  - 4′,4‴-diphenyl-2,2′:6′,2″:6″,2‴:6‴,2⁗-quinquepyridyl-, tetraphenylboronate tetrahydrate, 311
  - dithiocyanatodi[4-tolylmercury(II) thiocyanato]-, 711
  - dithiocyanato(Me_4-[16]-tetene)-, 377
  - germanotungstate$(HCoGeW_{11}O_{40})^{7-}$, potassium salt hydrate, 283
  - heteropolytungstate, 283
  - hexaaquo-, chloride, 833
  - histidine pyridoxamine complexes, electronic spectra, formation constants and potentiometric titrations, 791
  - histidine(pyridoxamine)-, 791
  - macrocyclic tetramine complexes, synthesis, electronic and IR spectra and magnetic properties, 377
  - nitroacetate complexes, synthesis,electronic and IR spectra and magnetic properties, 247
  - nitroacetatobis(1,10-phenanthroline)-, tetrahydrate, 247
  - polypyridyl complexes, synthesis, IR spectra, electric conductance, magnetic properties and ligand reactivity, 21
  - pyridoxamine-, 791
  - tetrachloro-, methyltrioctylammonium and trioctylammonium salts, 623
  - thiocyanate complexes with mercury(II), synthesis, electronic and IR spectra, electric conductance, magnetic properties and "softness", 711
- Cobalt(II) (cont)
  - tris(2,2′-bipyridyl)-, dithiocyanato-1-naphthylmercurate(II), 711
  - tris(2,2′-bipyridyl)-, dithiocyanato-4-tolylmercurate(II), 711
  - tris(1-phenyl-3-methyl-4-benzoyl-pyrazol-5-onato)-, trioctylammonium salt, 461
- Cobalt(III)
  - binuclear μ-peroxo complexes, synthesis, electronic, IR and NMR(^{1}H) spectra, electric conductance, magnetic properties, thermogravimetric analysis and ligand reactivity, 21
  - bis[8-bromodecahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - bis[8-chlorodecahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511

  - bis[9-chlorodecahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - bis[8,9-dichloroenneahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - bis[9,12-dichloroenneahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - bis(di-2-pyridylketone)-μ-peroxobis(2,2′,2″- terpyridyl)di-, perchlorate, 21
  - bisethylenediamine(2,4-pentanedionato)-, NMR (^{1}H,^{13}C) spectra and base hydrolysis, 203
  - bis[8-iododecahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - bis[8,9,12-tribromooctahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - bis[8,9,12-trichlorooctahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - bis[8,9,12-triiodooctahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - bis[undecahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - [8-bromodecahydrodicarba-*nido*-undecaborate(2)][undecahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - carbaborane complexes, synthesis, electronic, IR and NMR (^{1}H,^{11}B) spectra, electric conductance, gel and HPLC chromatography, 511
  - carboxylate complex, 149
  - chloride, hydrated complex with poly(iminoethylene)-dithiocarbamate, 225
  - [8-chlorodecahydrodicarba-*nido*-undecaborate(2)][8,9-dichloroenneahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - [8-chlorodecahydrodicarba-*nido*-undecaborate(2)][undecahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - *trans*-dichlorotetrapyridine-, hydrogen dinitrate, 525
  - diketonate complex, base hydrolysis kinetics, 203
  - hexammine-, pentachlorocuprate(II), 365
  - [8-iododecahydrodicarba-*nido*-undecaborate(2)][undecahydrodicarba-*nido*-undecaborate(2)]- caesium salt, 511
  - tetrakis(di-2-pyridylketone)-μ-peroxobispyridinedi-, perchlorate tetrahydrate, 21
  - trismalonato-, oxidation of iron(II), kinetic study, 149

Cobaltacyclopentadienes, 743, 757

Cobaltacyclopentane, 748

Cobaltacyclopentenes, 745, 758

Cobaltacyclopropenes, 759

Cobinamides, NMR (^{1}H) spectra, spin-lattice relaxation, conformation changes and TLC, 321

Colour change, uranium(VI) complex, 483

Cone angles, carbaborane and cyclopentadienyl complexes, 663

Conformational analysis, molybdenum(II) square pyramidal diastereoisomers, 233

Conformation changes, cobalamins and cobinamides, 321

Controlled potential electrolysis, cobalt(II) and nickel(II) quinquepyridyl complexes, 311

Copper
  - reaction with ammonia and morpholine, 437
  - reaction with hydrogen peroxide and morpholine, 437

Copper(I)
  - acetoxytrihydroboratotrisdiphenyl(methyl)phosphine-, 299
  - *N*-aminorhodaninedibromodi- acetic acid solvate, 633
  - *N*-aminorhodanineiodo- *N*,*N*-dimethylformamide solvate, 633
  - bromo-*N*-methylrhodanine-, 633
  - chloro-*N*-aminorhodanine- acetic acid solvate, 633
  - chloro-*N*-methylrhodanine-, 633
  - chlorotrisdiphenyl(methyl)phosphine-, 299
  - di(*N*-aminorhodanine)- perchlorate acetic acid solvate, 633
  - hydroborate phosphine complex, synthesis and IR and NMR(^{1}H,^{11}B) spectra, 299
  - iodide, 611
  - iodo-*N*-methylrhodanine-, 633
  - penta(*N*-aminorhodanine)tetrabromotetra-, *N*,*N*-dimethylformamide solvate, 633
  - penta(*N*-aminorhodanine)tetrachlorotetra-, *N*,*N*-dimethylformamide solvate, 633
  - rhodanine complexes, synthesis, electric conductance and IR spectra, 633

- tetraacetonitrile-, hexafluorophosphate, 467
- tetra(N-methylrhodanine)-, perchlorate, 633
- trisdiphenyl(methyl)phosphineperchlorato-, 299

Copper(2+), reduction by trimethylphosphite in acetonitrile, 467

Copper(II)

- acetonitrilepentakistrimethylphosphite-, 467
- acetonitrile trimethylphosphite complex, formation, electronic spectrum, cyclic voltammetry and kinetic study, 467
- aminotrihydroxymethane complex, 361
- aquo-2,2′-bipyridylnitroacetato-, monohydrate, 247
- aquonitroacetato-1,10-phenanthroline-, monohydrate, 247
- bisacetonitriletetrakistrimethylphosphite-, 467
- bis(N-benzothiazolyl-N′-ethylthiourea)-, 557
- bis(N-benzothiazolyl-N′-methylthiourea)-, 557
- bis(camphorquinone dioxime monoanion)-, 503
- bis(camphorquinone dioxime monoanion)(camphorquinone dioxime dianion)di-, 503
- bisnitroacetato-, potassium salt monohydrate, 247
- bispentamethylcyclopentadienyl)-, 663

Copper(II) (cont)

- bis[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
- bromide, 543
- N,N′-butylenebis(2,4-pentanedioneiminato)-, 785
- N,N′-butylenebis(5,5,5-trifluoro-2,4-pentanedioneiminato)-, 785
- carboxylate quinoline complex, synthesis, electronic, ESR and IR spectra, magnetic properties, DTA and thermogravimetric analysis, 679
- chloride dihydrate, 543
- chloride complex, neutron diffraction and X-ray crystal structure, 365
- diaquoditheophyllinato-, 143
- diaquotetraacetatodi-, 557
- 2,6-dibenzothiazol-2-ylpyridinedibromo- monohydrate, 543
- 2,6-dibenzothiazol-2-ylpyridinedichloro- monohydrate, 543
- dibromo(Me_4-[16]-tetene)-, 377
- dichloro(Me_4-[16]-tetene)-, 377
- dihistidine-, 791
- dihistidine(pyridoxamine)-, 791
- dimorpholinedinitrato-, 437
- dinitrato(Me_4-[16]-tetene)-, 377
- dinitratotetramorpholine-, 437
- dioxime complexes, formation, electronic and ESR spectra, isomerism, ligand isomerism, optical activity and solvent extraction, 503
- di(2-oxoacetophenone oxime)-, 607
- di(2-oxo-1-naphthaldoxime)-, 607
- dipapaverinetetrapropionatodi-, 679
- di(2-peonol oxime)-, 607
- dipyridoxamine-, 791
- di(resacetophenone oxime)-, 607
- ditheophyllinato-, 143
- dithiocyanato(Me_4-[16]-tetene)-, 377
- N,N′-ethylenebis(2,4-pentanedioneiminato)-, 785
- N,N′-ethylenebis(5,5,5-trifluoro-2,4-pentanedioneiminato)-, 785
- N,N′-ethylene[(2,4-pentanedioneiminato)(5,5,5-trifluoro-2,4-pentanedioneiminato)]-, 785
- glutamine complexes, formation constants, 371
- glutamine(histamine)-, 371
- guanosine complexes, formation constants, 153
- histamine complexes in blood plasma, formation constants, 371
- histamine(histidine)-, 371
- histamine(threonine)-, 371
- histidine complexes, formation constants, 371
- histidine-, 791
- histidine pyridoxamine complexes, electronic spectra, formation constants and potentiometric titrations, 791
- histidine(pyridoxamine)-, 791
- macrocyclic tetramine complexes, synthesis, electronic and IR spectra and magnetic properties, 377
- morpholine complexes, synthesis, electronic and IR spectra, electrochemistry, formation constants and thermogravimetric analysis, 437

Copper(II) (cont)
  nalidixate complexes, formation constants, 153
  nitroacetate complexes, synthesis, ESR and IR spectra, magnetic properties and X-ray crystal structure, 247
  oxime phenolate complexes, electric conductance, ESR spectra and magnetic properties, 607
  pentaacetonitrile-, hexafluorophosphate, 467
  pentachloro-, hexamminecobalt(III) salt, 365
  N,N′-propylenebis(2,4-pentanedioneiminato)-, 785
  N,N′-propylenebis(5,5,5-trifluoro-2,4-pentanedioneiminato)-, 785
  purine complexes, synthesis, magnetic properties, electronic, ESR and IR spectra and DTA study, 143
  pyridoxamine-, 791
  reaction with apotransferrin in the presence of bicarbonate and aminotrihydroxymethane, kinetic study, 361
  reaction with iodide, kinetic study, 611
  sulphate, hydrated complex with poly(iminoethylene)-dithiocarbamate, 225
  tetradentate ketoimine complexes, synthesis, HPLC and isomerism, 785
  tetramorpholine-, 437
  threonine complexes, formation constants, 371
  transferrin complex, 361
  triaquotrinitrato-, 437
  trimorpholine-, 437
Corrinoids, see cobalamins and cobinamides
Critical adsorption concentration, oxime sulphonic acid complex, 453
Critical micelle concentration, oxime sulphonic acid complex, 453
18-Crown-6, 409
Crown ether complex, nickel(II), 409
Crystal structure, see X-ray crystal structure
Cyanide complex
  cobalamins and cobinamides, 321
  molybdenum(IV,IV,VI), 834
  tungsten(V), 779
Cyanocobalamin, 321
1-Cyanomethylpyridinium chloride, synthesis and IR and NMR (^{1}H) spectra, 303
Cyclic voltammetry
  cobalt(II) quinquepyridyl complexes, 311
  copper(II) acetonitrile trimethylphosphite complexes, 467
  iron(II) pentacyano thioamide complexes, 429
  nickel(II) quinquepyridyl complexes, 311
Cyclohexene
  hydroformylation, 592
  hydrogenation, 584
Cyclometallation reactions, formation of metallacycles, 771
1,5-Cyclooctadiene complexes, rhodium(I), 339
Cyclooctene, hydroformylation, 592
Cyclopalladation reactions, 179
Cyclopentadienedithiocarboxylate complexes, tin(II) and (IV), 187
Cyclopentadienide, sodium salt, 417
Cyclopentadienyl complexes
  cobalt, 133, 833
  iron, 1
  molybdenum(II), 233, 617
  neodymium(III), 423
  nickel, 133, 833
  palladium(II), 539
  rhenium(I), 441
  rhodium(I), 809
  samarium(III), 423
  tungsten(IV), 629
  uranium(III), 417
  ytterbium(II) and (III), 423
Cyclophosphamide complexes, antimony(III) and rhodium(II), 637
Cysteine complexes, uranium(VI), 537
Cystine complexes, cadmium(II) and zinc(II), 153
Cytidine, reaction with chromium(VI), 497
5′-Cytidinemonophosphate, reaction with chromium(VI), 497

D

Decachlorodecaborane(10), 134

Density, osmium carbohydrate polymers, 335
Deuterionitric acid, 525
Deuterionitric acid (^{15}N), 525
Deuterium, substitution in germamanium and tin organo halides, 397
Dialkene complexes, rhodium(I), 339
2,2′-Diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate complexes, americium(III) and lanthanides, 277
6,6′′-Diamino-4,4′′-diphenyl-2,2′:6′,2′′-terpyridyl, synthesis and mass spectrum, 303
1,2-Diaminoethane complexes
  cobalt(III), 203
  rhodium(III), 659
1,5-Diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate complexes with americium(III) and lanthanides and protonation constants, 277
Diarsenogermanotungstate($HAs_2GeW_{17}O_{62}$)$^{7-}$, potassium salt hydrate, 283
Diarsenostannotungstate($HAs_2SnW_{17}O_{62}$)$^{7-}$, potassium salt hydrate, 283
Diastereoisomers, molybdenum(II) square pyramidal complexes, conformational analysis, 233
Dibenzo-(*f*,*n*)-2,4,10,12-tetramethyl-1,5,9,13-tetraazacyclohexadeca-[16]-1,3,9,11-tetraene, *see* Me_4-[16]-tetene
2,6-Dibenzothiazol-2ylpyridine, synthesis, metal complexes, IR spectra, 543
Dibenzoyltartaric acid, Pfeiffer effect with lanthanide complexes, 195
Diboron tetraiodide, 319
Dibromoborane, 701
Dibromo(methyl)borane, 701
Dibromo(phenyl)borane, 701
6,6′′-Dibromo-4′-phenyl-2,2′:6′,2′′-terpyridyl, synthesis and mass spectrum, 303
Di-*t*-butylamine complex, vanadium(0), 209
Dibutylphenacylphosphonate complexes with lanthanides, 257
Dibutylphenacylphosphonic acid complex, lanthanum(III), 257
Dibutylphosphoric acid, synergic effect in solvent extraction, 683
closo-2,3-Dicarba-1-(η^6-benzene)-1-ferranonaborane(11), 77
closo-2,3-Dicarba-1-(η^6-benzene)-1-ruthenanonaborane(11), 77
5,6-Dicarba-*nido*-decaborane(12), 822, 824
closo-2,3-Dicarba-1-(η^6-hexamethylbenzene)ferranonaborane(11), 77
closo-2,3-Dicarba-1-(η^6-mesitylene)ferranonaborane(11), 77
Dichloroborane, 701
6,6′′-Dichloro-4,4′′-diphenyl-2,2′:6′,2′′-terpyridyl, synthesis and mass spectrum, 303
8,9-Dichloroenneahydrodicarba-*nido*-undecaborate(2) complexes, cobalt(III), 511
9,12 -Dichloroenneahydrodicarba-*nido*-undecaborate(2) complex, cobalt(III), 511
Dichloro(methyl)borane, 701
2,5-Dichlorophenyl,mercury(II) compounds, 243
Dichloro(phenyl)borane, 701
1,3-Di(4-chlorophenyl)triazene complex, rhodium(I), 573
2,6-Dicinnamoylpyridine, synthesis and IR and mass spectra, 303
Dicyanocobalamin, 321
Dicyanocobinamide, 321
2-Dicyclohexylarsinobenzoate complexes, cadmium(II), mercury(II), zinc(II), 667
2-Dicyclohexylbenzoic acid complexes, mercury(II), 667
Dielectric constants, chloride complexes in nitrobenzene, 623
1-Diethylamino-4-allylethylaminopent-2-yne, 563
3-Diethylaminopropene, 563
3-Diethylaminopropyne, 563
2-Diethylarsinobenzoate complexes, cadmium(II), zinc(II), 667
2-Diethylarsinobenzoic acid complexes, mercury(II), 667
Diethyleneglycol, solvent extraction, 383
Diethyleneglycol-1,4-bis(dibutylphosphate) complexes, uranium(VI), 683
Diethylenetriamine, *see* 1,4,7-triazaheptane
Di-2-ethylhexyl phosphoric acid, use in formation constant determination, 239
5,8-Diethyl-7-hydroxy-6-dodecanone, complex with dinonylnaphthalenesulphonic acid, 453
Diethyl(methyl)borane, 701

Diethyl(phenyl)phosphine complex, rhenium(I), 441
Diethyl-2-(4-sulphophenyl)ethylphosphonate, sodium salt, 61
Diethyl tartrate, Pfeiffer effect with lanthanide complexes, 195
Differential thermal analysis, *see* DTA
Difluoroborane, 701
Difluoro(*n*-butyl)borane, 701
Difluoro(ethyl)borane, 701
Difluoro(methyl)borane, 701
1,1-Difluoro-4-phenyl-1,3-butanedionate, gallium(III) and indium(III) compounds, 9
Difluorophosphate complexes, ruthenium(II) and silver(I), 97
Difluoro(*i*-propyl)borane, 701
1,1-Difluoro-4-[2′-thienyl]-1,3-butanedionate, gallium(III) and indium(III) compounds, 9
2,2′-Difluorovinyl compounds, sulphur(VI), 723
Digermanotungstate($HGe_2W_{11}O_{40}$)$^{5-}$, potassium salt hydrate, 283
Di-Grignard reagents, for formation of metallacycles, 744
2,5-Dihydrofuran, hydroformylation, 599
3,4-Dihydroxybenzoate dianion complexes, aluminium(III), 313
β-Diketonate complexes
  chromium(III), 383
  cobalt(II), 461
  cobalt(III), 203
  gallium(III), 9
  indium(III), 9, 457
  nickel(II), 461
  uranium(VI), 483
  zinc(II), 387
Dilithio reagents, for formation of metallacycles, 744
1,2-Dimethoxyethane, 417
1,2-Dimethoxyethane complex, ytterbium(II), 423
Dimethoxy(methyl)phosphine oxide, 467
N,*N*-Dimethylacetamide
  solvent extraction, 383
  solvent for lanthanide ion complexation studies, 49
2-Dimethylarsinobenzoate complex, mercury(II), 667
Dimethyl(ethyl)borane, 701
N,*N*-Dimethylformamide
  electric conductance of solutions of lanthanide toluene-4-sulphonate complexes, 253
  solvates, 633
  solvent extraction with, 383
  solvolysis of chromium(III) complex, 191
3,7-Dimethyl-7-methoxyoctene, isomerisation, 580
3,7-Dimethylocta-1,6-diene, isomerisation, 579
2,3-Dimethylpentanal, 589
2,9-Dimethylphenanthroline complex, zinc(II), solvent extraction, 387
2,9-Dimethyl-4,7-diphenylphenanthroline complex, zinc(II), solvent extraction, 387
4,7-Dimethylphenanthroline complex, zinc(II), solvent extraction, 387
Dimethyl(phenyl)phosphine complex, tungsten(IV), 629
Dimethyl(phenyl)silane, 601
4,7-Diphenylphenanthroline complex, zinc(II), solvent extraction, 387
Dimethylphosphidomethyl complex, rhenium(III), 37
1,2-Dimethylselenoethane complex, platinum(II), 289
Dimethylsulphonium benzoylmethylide complex, palladium(II), 539
Dimethylsulphoxide
  complex, iron(II), 429
  electric conductance and solvolysis of lanthanide toluene-4-sulphonate complexes, 253
  NMR (^{1}H) spectra with lanthanide complexes, 253
  reaction with iron(II) pentacyano thioamide complex, 429
  solvent extraction with, 383
  solvolysis of chromium(III) complex, 191
Dimethylsulphoxide complexes
  cobalt(II), 711
  mercury(II), 819
  nickel(II), 711
Dimethyl tartrate, Pfeiffer effect with lanthanide complexes, 195
1,2-Dimethylthioethane complexes, palladium(II) and platinum(II), 289

1,2-Dimethylthioethene complexes, palladium(II) and platinum(II), 289
1,3-Dimethylthiopropane complexes, palladium(II) and platinum(II), 289
3,4-Dimethylthiotoluene complexes, palladium(II) and platinum(II), 289
Dinitrogen complexes
  magnesium, 421
  rhenium(I), 37, 441
Dinonylnaphthalenesulphonic acid
  complex with 5,8-diethyl-7-hydroxy-6-dodecanone, 453
  interfacial tension, spreading pressure and inverse micelle formation, 541
Diolefin, *see* dialkene
DIOP, rhodium(I) complex as hydroformylation catalyst, 599
Diphenolates, acid dissociation constants and formation constants of alumninium(III) complexes, 313
Diphenylacetylene, 549, 575, 761, 765
2-Diphenylarsinobenzoate complex, mercury(II), 667
Diphenyl(methyl)phosphine complex, copper(I), 299
Diphenyl[pentacarbonylchromium(0)]phosphido complexes, mercury(II), 819
Diphenyl[pentacarbonylmolybdenum(0)]phosphido complexes, mercury(II), 819
Diphenyl[pentacarbonyltungsten(0)]phosphido complexes, mercury(II), 819
Diphenylphosphide complex, rhodium(I), 578
1,2-Diphenylphosphinoethane complex, rhodium(I), 573
η^2-Diphenylphosphinophenyl complexes, osmium(III), 83
Diphenyl(2-pyridyl)phosphine complex, rhodium(I), 572
4,4‴-Diphenyl-2,2′:6′,2″:6″,2‴:6‴,2⁗-quinquepyridyl complexes, cadmium(II), cobalt(I) and (II), nickel(I) and (II), 311
4,4″-Diphenyl-2,2′:6′,2″-terpyridyl-6(1H),6″(1″H)-dione, synthesis, 303
1,3-Diphenyltriazene complex, rhodium(I), 573
DIPHOL, rhodium(III) complex as hydroformylation catalyst, 599
Diphosphogermanotungstate$(HP_2GeW_{17}O_{62})^{7-}$, potassium salt hydrate, 283
Diphosphostannotungstate$(HP_2SnW_{17}O_{62})^{7-}$, potassium salt hydrate, 283
Dipicolinic acid, *see* pyridine-2,6-dicarboxylic acid
N,*N*′-Dipropionato-1,2-diaminoethane complex, rhodium(III), 659
Di-*i*-propyl(methyl)borane, 701
Di-*n*-propyl(methyl)borane, 701
2,2′-Dipyridylamine complex, vanadium(0), 209
2,2′-Dipyridyldisulphide complex, vanadium(0), 209
Di-2-pyridylketone complexes, cobalt(II) and(III), synthesis and reactivity, 21
N,*N*′-Disalicylidene-3,4-phenylenediamine-1-ethylbenzoate complex, zirconium(IV), 487
Disproportionation, lanthanide(II) halides and oxides, 219
Dissociation enthalpies, caesium trihalides, 215
Dithiocarboxylate complexes, tin(II) and (IV), 187
Dithiooxamide complex, iron(II), 429
2-Di(4-tolyl)arsinobenzoate complex, mercury(II), 667
Di-(4-tolyl)tartaric acid, Pfeiffer effect with lanthanide complexes, 195
1,3-Di(4-tolyl)triazene complex, rhodium(I), 573
Dodecahedral cluster compounds, 133
Dodecahydrodicarba-*nido*-decaborate(2), 663
Dodecane, in solvent extraction, 383
DTA
  copper(LI) carboxylate complex, 679
  copper(II) purine complexes, 143
  magnesium azide hydrazine complex, 421
  molybdenum(V) oxo oxoquinoline complexes, 163
  molybdenum(VI) and tungsten (VI) oxides with sodium chloride, 479
  zirconium(IV) sulphophosphonates, 61
Dysprosium(III)
  complexes with 2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate, 277
  toluene-4-sulphonate, 253

E

- EDTA, *see* ethylenediaminetetraacetate
- EHMO calculation, mercury(II) alkyl, 413
- Eight-coordination, zirconium(IV), 487
- Electric conductance
  - antimony(III) and bismuth(III) halide thiazole complexes, 655
  - biacetyldihydrazone complexes, 831
  - cadmium(II)
    - halide mercaptide complexes, 157
    - phosphine selenide and phosphine sulphide complexes, 781
  - chloride complexes, 623
  - cobalt(II)
    - polypyridyl complexes, 21
    - thiocyanate complexes, 711
  - cobalt(III)
    - binuclear complexes, 21
    - carbaborane complexes, 511
  - copper(I) rhodanine complexes, 633
  - copper(II) oxime complexes, 607
  - mercury(II)
    - halide mercaptide complexes, 157
    - phosphine selenide and phosphine sulphide complexes, 781
    - thiocyanate complexes, 711
  - molybdenum(V) oxo oxoquinoline complexes, 163
  - nickel(II)
    - macrocyclic tetramine complexes, 377
    - thiocyanate complexes, 711
  - organotellurium carboxylates, 13
  - osmium(IV) chloride complexes, 83
  - phosphonium salts, 83
  - rhenium(I)
    - alkylidyne isocyanide complexes, 803
    - dinitrogen phosphine complexes, 441
  - rhenium(III)
    - chloride hydride phosphine complex, 441
    - iodide phenylamide phosphine complex, 441
  - rhenium(V)
    - amido and imido complexes, 31
    - hydride phosphine complex, 441
  - rhodium(I) dialkene complexes, 339
- Electric conductance (cont)
  - ruthenium(II) phosphine complexes, 97
  - ruthenium(II)-silver(I) bonded complex, 97
  - silver(I) rhodanine complexes, 633
  - thiazolylthiourea complexes, 557
  - zinc(II)
    - halide mercaptide complexes, 157
    - phosphine selenide and phosphine sulphide complexes, 781
- Electrochemistry
  - copper(II) morpholine complexes, 437
  - heteropolytungstates, 283
  - iron(II) pentacyano thioamide complexes, 429
  - osmium(V) and (VIII) in aqueous solution, 175
- Electronic spectra
  - aluminium(III) methylthymol blue complexes, 405
  - bipyrimidine complexes, 53
  - chromium(II) binuclear carboxylate, 135
  - chromium(III) thiocyanate complex, 191
  - cobalt(II)
    - histidine pyridoxamine complexes, 791
    - macrocyclic tetramine complexes, 377
    - thiocyanate complexes, 711
  - cobalt(III)
    - binuclear complexes, 21
    - carbaborane complexes, 511
    - polyimine dithiocarbamate complex, 225
  - copper(II)
    - acetonitrile trimethylphosphite complexes, 467
    - carboxylate complex, 679
    - dioxime complexes, 503
    - histidine pyridoxamine complexes, 791
    - macrocyclic tetramine complexes, 377

    morpholine nitrate complexes, 437
    oxime complexes, 607
    polyimine dithiocarbamate complex, 225
    purine complexes, 143
  cyanide iron(II) complexes, 429
  heteropolytungstates, 283
  iron(II)
    carbaborane and cyclopentadienyl complexes, 663
    pentacyano thioamide complexes, 429
    polyimine dithiocarbamate complex, 225
  iron(III)
    fluoro porphyrin complex, 737
    polyimine dithiocarbamate complex, 225
  lanthanide toluene-4-sulphonate complexes, 253
  lead(II) bromide complexes, 169
  magnesium amide dinitrogen complex, 421
  manganese(II)
    biacetyldihydrazone complex, 831
    bipyridyl and bipyridyl dioxide complexes, 327
  methylthymol blue aluminium(III) complexes, 405
  molybdenum(IV,IV,VI) cyanide oxo complex, 834
Electronic spectra (cont)
  molybdenum(V) polymeric aquo oxo ion, 109
  nickel(II)
    histidine pyridoxamine complexes, 791
    macrocyclic tetramine complexes, 377
    polyimine dithiocarbamate complex, 225
    thiocyanate complexes, 711
  osmium(VI) and (VIII) in aqueous solution, 175
  poly(iminoethylene)dithiocarbamate and its complexes, 225
  rhodium(II) carboxylate cyclophosphoramide complexes, 637
  rhodium(III) aminopolycarboxylate diamine complex, 659
  thioamides and their pentacyanoiron(II) complexes, 429
  tin organometallic compounds, 1
  uranium(III) cyclopentadienyls, 417
  uranium(VI) dioxo fluoride complex, 835
  vanadium(IV) oxo polyimine dithiocarbamate complex, 225
  zirconium(IV) tetradentate Schiff base complex, 487
  see also photoacoustic spectra
Electron microscopy, silver(II) oxide, 183
Electron probe microanalysis, uranium apatites, 343
Electron spin resonance, *see* ESR
Emission spectra, *see* fluorescence spectra
Enantiomers, organotin compounds, 1
Enneabromoenneaborane(9), 134
Enneaiodoenneaborane(9), 319
Enthalpy of formation
  boron compounds, 701
  lanthanide(II) halides and oxides, 219
  lanthanide 3+ hydrated ions, 219
1,2-Epoxycyclohexan-3-ol, isomerisation, 581
Erbium(III)
  complexes with 2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N*′,*N*′- tetraacetate, 277
  toluene-4-sulphonate, 253
Erratum, caesium perbromate crystal structure, 567
ESR spectra
  bipyrimidine anion complexes, 53
  chromium(V) ribonucleotide complexes, 497
  cobalt(II) quinquepyridyl complexes, 311
  copper(II)
    carboxylate complex, 679
    dioxime complex, 503
    nitroacetate complex, 247
    oxime complexes, 607
    polyimine dithiocarbamate complex, 225
    purine complexes, 143
  iron(III)
    fluoro porphyrin complexes, 737
    polyimine dithiocarbamate complex, 225
  manganese(II)
    biacetyldihydrazone complex, 831

    bipyridyl complexes, 327
ESR spectra (cont)
  nickel(I) quinquepyridyl complexes, 311
  nickel(II), polyimine dithiocarbamate complex, 225
  osmium(III) complexes, 83
  quinquepyridyl complexes, 311
  rhenium(IV) imide complex, 37
  tungsten(V) phenoxo complex, 641
  vanadium(IV) oxo polyimine dithiocarbamate complex, 225
Ethambutol complex, copper(II), 153
Ethanol
  photochemical decomposition, 839
  solvates of mercury(II) complexes, 667
  solvation of zirconium(IV) complex, 487
  solvolysis of chromium(III) complex, 191
Ethene, *see* ethylene
1-Ethoxycarbonylmethylpyridinium bromide, synthesis and IR and NMR (^{1}H) spectra, 303
Ethyl compounds
  phosphorus(V), 123
  rhenium(I), 803
Ethyl acrylate, hydrogenation, 584
Ethyl-4-amino-3-nitrobenzoate, 487
Ethyl-3,4-diaminobenzoate, 487
Ethylene
  carbonylation, 601
  deuterium substitution, 579
Ethylene complex, platinum(II), 57
N,*N*′-Ethylenebis(2,4-pentanedionimine) complexes, copper(II), nickel(II), palladium(II), 785
N,*N*′-Ethylenebis(5,5,5-trifluoro-2,4-pentanedioneimine) complexes, copper(II), nickel(II), palladium(II), 785
Ethylenediamine, *see* 1,2-diaminoethane
Ethylenediaminediacetate complex, cadmium(II), 153
Ethylenediaminedipropionate complex, rhodium(III), 659
Ethylenediaminetatraacetate complex, cadmium(II), 153
[(Ethylenedioxy)diethylenedinitrilo]tetraacetate complexes, thorium(IV) and uranium(VI), 827
Ethyleneglycol, in solvent extraction, 383
Ethyleneglycol-1,2-bis(dibutylphosphate) complexes, uranium(VI), 683
N,*N*′-Ethylene[(2,4-pentanedioneimine)(5,5,5-trifluoro-2,4-pentanedioneimine)] complexes, copper(II), nickel(II), palladium(II), 785
2-Ethylhexanol, 585
Ethylidyne complex, iron(0), 842
Ethyl(methyl)-*n*-propylborane, 701
2-Ethylpyridine complexes, rhodium(I), 339
Ethyne, hydrosilylation, 601
Europium(II) fluoride and oxide, improbability of disproportionation, 219
Europium(III)
  aminopolycarboxylate complexes, reaction with ytterbium(3+), kinetic study, 707
  bis(trimethylphosphine oxide)tris(bistrimethylsilylamido)-, 307
Europium(III) (cont)
  chiral complex, fluorescence and circularly polarised fluorescence spectra, Pfeiffer effect and formation constants, 195
  complexes with 2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate, 277
  cyclohexanediaminetetraacetato-, 707
  dihydroxofluoro-, 239
  ethylenediaminetetraacetato-, 707
  propylenediaminetetraacetato-, 707
  toluene-4-sulphonate, 253
  trimethylphosphine oxide tris(bistrimethylsilylamido)-, 307
  tris(dibutylphenacylphosphonato)-, 257
  tris(pyridine-2,6-dicarboxylato)-, 195
  tris(pyridine-2,6-dicarboxylato)-, dibenzoyltartaric acid complex, 195
  tris(pyridine-2,6-dicarboxylato)-, diethyl tartrate complex, 195
  tris(pyridine-2,6-dicarboxylato)-, dimethyl tartrate complex, 195
  tris(pyridine-2,6-dicarboxylato)-, di-(4-tolyl)- tartaric acid complex,

Extended Hueckel Molecular Orbital method, *see* EHMO

F

Factor group analysis
  chromium(III) chloride, 475
  nickel molybdate, 331
Fenske - Hall calculations, boranes and ferraboranes, 521
Ferraboranes, molecular orbital calculation, 521
Ferracyclobutane, 742
Ferracyclopentadienes, 750, 766
Ferracyclopentanes, 743, 751, 766
Ferracyclopentenes, 766
Fluorescence spectra
  lanthanide chiral complexes, 195
  terbium(III) hydroxycarboxylate salicylate complexes, 693
  uranium(VI), 343, 835
Fluorides and fluoro-complexes
  cerium(III), 239
  europium(III), 239
  iron(III), 737
  organotelluronium salt, 13
  rhenium(V), 31
  ruthenium(II), 97
  scandium(III), 735
  sulphur(VI), 723
  terbium(III), 239
  uranium(VI), 835
  ytterbium(III), 239
Fluorite-bearing system, lanthanide complexes, 239
Fluoroalkyl compounds, silver(I) and sulphur(VI), 723
Fluoroapatite, uranium(VI) in, 343
Fluoroborane, 701
Fluoro(di-*n*-butyl)borane, 701
Fluoro(dimethyl)borane, 701
Fluorohydroxycarbonateapatite, uranium(VI) in, 343
1-[4′-Fluorophenyl]-4,4,4-trifluoro-1,3-butanedionate, gallium(III) and indium(III) compounds, 9
Fluoropolytungstates, synthesis, polarography, hydrolysis kinetics and reactions with Al^{3+} and Co^{3+}, 263
Fluorotungstate $(HF_2W_{12}O_{38})^{5-}$, 263
Fluorotungstate $(H_2FW_{12}O_{39})^{5-}$, 263
Fluorotungstate $(HF_3W_{12}O_{37})^{4-}$, acid and potassium salt, 263
Fluorotungstate $(H_2F_2W_{12}O_{38})^{4-}$, acid and potassium salt, 263
Fluxional behaviour
  gallium(III) and indium(III) β-diketonates, 9
  palladium(II) and platinum(II) selenoether and thioether complexes, 289
Force constants
  germanium and tin organo halides, 397
  uranium apatites, 343
Formamide, solvent extraction of chromium(III) diketonates, 383
Formation constants
  aluminium(III) diphenolate complexes, 313
  aluminium(III) methylthymol blue complexes, 405
Formation constants (cont)
  cadmium(II) complexes, 153
  cobalt(II) β-diketonate complexes, 461
  cobalt(II) histidine pyridoxamine complexes, 791
  copper(II)
    guanosine complexes, 153
    histamine complexes in blood plasma, 371
    histidine pyridoxamine complexes, 791
    morpholine complexes, 437
    nalidixate complexes, 153
  determination by NMR (^{13}C) spectra, 145
  determination by partition between di-2-ethylhexylphosphoric acid and water, 239
  indium(III) diketonate complexes, 457
  iron(III) complexes with pyridine carboxylic acids, 269
  lanthanide
    chiral complexes complexed by tartaric acid and derivatives, 195

    - diaminotetracarboxylate complexes, 277
    - fluoride and hydroxide complexes, 239
    - halide and nitrate complexes,49
  - lead(II) mercaptoethanol complexes, 836
  - nickel(II) β-diketonate complexes, 461
  - nickel(II) histidine pyridoxamine complexes, 791
  - potentiometric determination by competition method for systems with an insoluble component, 153
  - silver(I) - acetonitrile complexes, 145
  - terbium(III) hydroxycarboxylate salicylate complexes, 693
  - thorium(IV) aminopolycarboxylate complexes, 827
  - uranium(VI) aminoacid complexes, 537
  - uranium(VI) aminopolycarboxylate complexes, 827
  - zinc(II) aminothiazoline complexes, 153
  - zinc(II) histidine pyridoxamine complexes, 791

G

- Gadolinium(III)
  - complexes with 2,2′-diaminoether,*N*,*N*,*N*′,*N*′-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate, 277
  - trimethylphosphine oxide tris(bistrimethylsilylamido)-, 307
  - tris(dibutylphenacylphosphonato)-, 257
- Gallium(III)
  - β-diketonates, synthesis,NMR (^{13}C,^{19}F) spectra and fluxional behaviour, 9
  - in heteropolytungstates, 283
  - tetrachloro-, methyltrioctylammonium and trioctylammonium salts, 623
  - tris(1,1-difluoro,4-phenyl-1,3-butanedionato)-, 9
  - tris(1,1-difluoro,4-[2′-thienyl]-1,3-butanedionato)-, 9
  - tris(1-[4′-fluorophenyl]-4,4,4-trifluoro-1,3-butanedionato)-, 9
  - tris(1-[4′-methylphenyl]-4,4,4-trifluoro-1,3-butanedionato)-, 9
  - tris(1-[2′-naphthyl]-4,4,4-trifluoro-1,3-butanedionato)-, 9
  - tris(1-phenyl-4,4,4-trifluoro-1,3-butanedionato)-, 9
- Gallium(III) (cont)
  - tris(1-[2′-thienyl]-4,4,4-trifluoro-1,3-butanedionato)-, 9
  - tris(1,1,1-trifluoro-2,4-pentanedionato)-, 9
- Gallogermanotungstate ($HGaGeW_{11}O_{40})^{6-}$, potassium salt hydrate, 283
- Gallostannotungstate ($HGaSnW_{11}O_{40})^{6-}$, potassium salt hydrate, 283
- Gel chromatography, cobalt(III) carbaborane complexes, 511
- Gel electrophoresis, osmium carbohydrate polymers, 335
- Gel filtration, osmium carbohydrate polymers, 335
- Geochemical studies, lanthanides in fluorite-bearing system, 239
- Germanium
  - bromotrimethyl-, 397
  - chlorotrimethyl, 397
  - in heteropolytungstates, 283
  - organohalides, isotopic substitution, IR and Raman spectra, normal coordinate analysis, force constants and bond lengths, 397
- Germanosilicotungstate ($HGeSiW_{11}O_{40})^{5-}$, potassium salt hydrate, 283
- Germanostannotungstate ($HGeSnW_{11}O_{40})^{5-}$, potassium salt hydrate, 283
- Germanozinctungstate ($HGeZnW_{11}O_{40})^{7-}$, potassium salt hydrate, 283
- Gluconate compound, osmium polymer, 335
- Glucose compound, osmium polymer, 335
- Glutamine complexes, copper(II), 371
- Gold(I)
  - (dodecacarbonyldihydridotetraosmium)bistriethylphosphinedi-, 105
  - (dodecacarbonyldihydridotetraosmium)bistriphenylphosphinedi-, 105
  - (dodecacarbonyltrihydridotetraosmium)triethylphosphine-, 105
  - (dodecacarbonyltrihydridotetraosmium)triphenylphosphine-, 105
  - phosphine complexes with osmium cluster carbonyl hydrides, synthesis, IR and mass spectra, X-ray crystal structure, isomerism and reactions, 105
- Guanosine, reaction with chromium(VI), 497
- Guanosine complexes, copper(II), 153
- 5′-Guanosinemonophosphate, reaction with chromium(VI), 497

H

- Hafnacyclopentane, 749
- Hafnium(0)

  tris(2,2′-bipyridyl)-, synthesis, 209
  tris(1,10-phenanthroline)-, synthesis, 209
Hafnium(IV)
  bis[dodecahydrodicarba-*nido*-dodecaborate(2)]-, 663
  di(η^1-cyclopentadienyl)di(η^5-cyclopentadienyl)-, 663
  tetrachloro-, 209
Halogenation of carbaboranes induced by γ-radiation, 511
HDNNS, *see* dinonylnaphthalenesulphonic acid
Heptachloroenneaborane(9), 134
Heptafluoroethoxymethylamine, 129
Heptafluoro-*N*-methylacetamide, 129
Heptanal, 591
Heptapolyphosphate, magnesium and tetramethylammonium salts, 69
Hept-1-ene, isomerisation, 579
Heteropolytungstates
  syntheses, electronic spectra, electrochemistry and *X*-ray diffraction, 283
  see also fluoropolytungstates
Hexachloroennaborane(9), 134
Hexafluoroformamidine, 129
Hexafluoro-1-methyldiaziridine, 129
1,1,1,5,5,5-Hexafluoropentane-2,4-dionate complexes
  rhodium(I), 573
  uranium(VI), 483
3,3′,4,4′,5,5′-Hexahydroxydiphenylsulphone dianion complexes, aluminium(III), 313
Hexamethoxydiphosphine cation, 467
Hexamethylbenzene complex, iron(II), 77
Hexamethylphosphoramide complex, indium(III), 355
Hexane, formation from hex-1-ene, 839
Hexaphenyltriphosphazine, 535
Hexapolyphosphate, magnesium and tetramethylammonium salts, 69
Hex-1-ene, 592
Hex-3-yne, formation of metallacycles, 755
High pressure liquid chromatography, *see* HPLC
Histamine complexes, copper(II), 371
Histidine complexes
  cobalt(II), 791
  copper(II), 371, 791
  nickel(II) and zinc(II), 791
Holmium(III)
  chiral complex, fluorescence and circularly polarised fluorescence spectra, Pfeiffer effect and formation constant, 195
  complexes with 2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate, 277
  toluene-4-sulphonate, 253
  tris(pyridine-2,6-dicarboxylato)-, 195
HPLC
  ammonolysis of trimetaphosphate, 425
HPLC (cont)
  cobalt(III) carbaborane complexes, 511
  nickel(II) and palladium(II) tetradentate ketoimine complexes, 785
Hydrazine, 569
Hydrazine complexes, magnesium, 421
Hydrazinium salt, 735
Hydride, displacement from rhodium(I) complex, 573
Hydrides and hydrido- complexes
  boron, 134
  iron(0), 842
  osmium(III), 83
  osmium cluster carbonyl, 105
  rhenium(I), 441
  rhenium(III), 37, 441
  rhenium(V), 441
  rhodium(I), 569, 839
  tungstenIV), 629
Hydroborate, 570
Hydroborate complex, copper(I), 299
Hydro-1,5-diazabicyclo[5,4,0]-undec-1-ene, trichloro(triethylphosphine)platinate(II)salt, 89
Hydrogen, 441, 487, 591
Hydrogen bonding, uranium(VI) ammonia diketonate oxo complex, 483

Hydrogen dinitrate salts, synthesis, IR and Raman spectra and X-ray crystal structure and diffraction, 525
Hydrogen Peroxide
  reaction with copper and ammonia or morpholine, 437
  reduction of and catalytic decomposition by osmium(VIII), 175
  synthesis of vanadium(V) peroxocomplexes, 553
Hydroformylaion
  catalysis by rhodium(I) complex, 584
  catalyst regeneration, 596
  chiral, 597
  selectivity, 590
Hydrogenation, catalysis by rhodium(I) complex, 581
Hydrolysis
  chromium(III) thiocyanate complex, 191
  fluoropolytungstates, 263
  hydro(pyrrolyl)borates, 673
Hydroperoxides, reaction with rhodium(I) complexes, 584
Hydroxide complexes
  aluminium(III), 313
  cerium(III), 239
  europium(III), 239
  iron(III), 799
  molybdenum(V), 163
  platinum(IV), 57
  scandium(III), 735
  terbium(III), 239
  ytterbium(III), 239
Hydroxyapatite, uranium(VI) in, 343
Hydroxycarboxylate complexes, terbium(III), 693
Hyponitrite complex, rhodium(III), 577

I

Imido complexes
  rhenium(IV), 37
  rhenium(V), 31, 37
  tungsten(VI), 561
Iminodiacetate complexes, thorium(IV) and uranium(VI), 827
Indium(III)
  bis(4-benzoylpyrazol-5-onato)dichloro-, methyltrioctylammonium and trioctylammonium salts, 457
  bis(hexamethylphosphoramide)trichloro-, 355
  bis(4-lauroylpyrazol-5-onato)dichloro-, trioctylmethylammonium salt, 457
  bis(1-phenyl-3-methyl-4-benzoylpyrazol-5-onato)chloro-, 457
  bis(1-phenyl-3-methyl-4-lauroylpyrazol-5-onato)chloro-, 457
  bis(1-phenyl-3-methyl-4-[2-thenoyl]-pyrazol-5-onato)chloro-, 457
  bis[4-(2-thenoyl)pyrazol-5-onato]dichloro-, trioctylmethylammonium salt, 457
  bis(tetraphenylbuta-1,3-diyl)-, tetraphenylarsonium salt and maleic anhydride adduct, 549
  chloride β-diketonate complexes, formation constants,solvent extraction, synergic effects, 457
  chloride phosphine oxide complex, synthesis, IR and Raman spectra and X-ray crystal structure, 355
  chloro(pyridine)(tetraphenylbuta-1,3-diene-1,4-diyl)-, 549
  β-diketonates, synthesis, NMR $(^{13}C,^{19}F)$ spectra and fluxional behaviour, 9
  organometallic compounds, synthesis and NMR $(^{1}H,^{13}C)$ spectra, 549
  tetrachloro-, methyltrioctylammonium and trioctylammonium salts, 623
  tetrachloro-, tetraphenylarsonium salt, 549
  trichlorotripyridine-, 549
  trimethylphosphine oxide tris(bistrimethylsilylamido)-, 307
  tris(1,1-difluoro-4-phenyl-1,3-butanedionato)-, 9
  tris(1,1-difluoro-4-[2′-thienyl]-1,3-butanedionato)-, 9
  tris(1-[4′-fluorophenyl]-4,4,4-trifluoro-1,3-butanedionato)-, 9
  tris(1-[4′-methylphenyl]-4,4,4-trifluoro-1,3-butanedionato)-, 9
  tris(1-[2′-naphthyl]-4,4,4-trifluoro-1,3-butanedionato)-, 9
  tris(1-phenyl-3-methyl-4-benzoylpyrazol-5-onato)-, 457
  tris(1-phenyl-3-methyl-4-lauroylpyrazol-5-onato)-, 457
  tris(1-phenyl-3-methyl-4-[2-thenoyl]-pyrazol-5-onato)-, 457
  tris(1-phenyl-4,4,4-trifluoro-1,3-butanedionato)-, 9

  tris(1-[2′-thienyl]-4,4,4-trifluoro-1,3-butanedionato)-, 9
  tris(1,1,1-trifluoro-2,4-pentanedionato)-, 9
Inosine, reaction with chromium(VI), 497
5′-Inosinemonophosphate, reaction with chromium(VI), 497
Interfacial tension
  dinonylnaphthalenesulphonic acid, 541
  oxime sulphonic acid complex, 453
Inverse micelle formation, dinonylnaphthalenesulphonic acid, 541
Iodide, reactions with
  copper(II), kinetic study, 611
  platinum(II) complex, 201
Iodides and iodo complexes
  antimony(III) and bismuth(III), 655
  cadmium(II), 137, 157, 781, 826
  cobalt(II), 543
  copper(I), 633
  mercury(II), 137, 157, 667, 781
  palladium(II), 179, 289
  platinum(0), 89
  platinum(II), 289
  rhenium(III), 441
  tin(II), 826
  tungsten(IV), 629
  vanadium(II), 209
  zinc(II), 157, 781
Iodine, 441
Iodine clock, 611
Iodine(I),
  bromoiodo-, caesium salt, 215
  dibromo-, caesium salt, 215
  dichloro-, caesium salt, 215
7-Iodo-5,6-dicarba-*nido*-decaborane(12), 822
8-Iodododecahydrodicarba-*nido*-undecaborate(2) complexes, cobalt(III), 511
Iodomethane, 441
Ion exchange, *see* anion exchange, cation exchange
IR spectra
  acetoxyhydroborate complex, 299
  acyl complex, 83
  alkylidyne complexes, 803
  aluminium(III) chloride complexes, 393
  amide complexes, 31, 421
  amine complexes, 633
  ammonium derivatives, 535
  antimony(III) halide thiazole complexes, 655
  arene complexes, 77
  arylphosphines, 535
  azide complex, 421
  biacetyldihydrazone complexes, 831
  bipyridyl complexes, 711
  bipyrimidine complexes, 53
  bismuth(III) halide thiazole complexes, 655
  bromide complexes, 157, 179, 377, 397, 655
  carbaborane complexes, 511
  carbon dioxide complex, 441
  carbonyl complexes, 45, 53, 97, 105, 209, 233, 441, 570, 663, 689
  carboxylate complexes, 31, 135, 667, 679
  carboxylic acids and their mercury(II) complexes, 667
  chloride complexes, 83, 89, 157, 179, 243, 355, 377, 393, 397, 475, 655, 689
  chromium(II) binuclear carboxylate thiocyanate complex, 135
IR spectra (cont)
  chromium(III) chloride, 475
  cobalt(-I) carbonyl complexes, 45
  cobalt(II) macrocyclic complexes, 377
  cobalt(II) nitroacetate complexes, 247
  cobalt(II) polypyridyl complexes, 21
  cobalt(III) binuclear μ-peroxo complexes, 21
  cobalt(II) carbaborane complexes, 511
  copper(I) acetoxytrihydroborate complex, 299
  copper(I) halide complexes, 633
  copper(II) carboxylate complex, 679
  copper(II) macrocyclic complexes, 377

  copper(II) morpholine nitrate complexes, 437
  copper(II) nitroacetate complexes, 247
  crown ether complexes, 409
  cyanide complex, 779
  cyclophosphamide complexes, 637
  difluorophosphate complex, 97
  dimethylsulphoxide complexes, 711
  dinitrogen complex, 421, 441
  fluoride complexes, 553, 735, 737
  fluoroalkyl compounds, 723
  fluorothio compounds, 723
  germanium organo halides, 397
  hexamethylphosphoramide complex, 355
  hydrates, 543, 667
  hydrazine complexes, 421, 735
  hydrazinium salt, 735
  hydrazone complexes, 179
  hydride complexes, 441, 570, 629
  hydroborate complex, 299
  hydrogen dinitrate, 525
  hydroxide complexes, 163, 735, 799
  imide complexes, 31
  indium(III) chloride phosphine oxide complex, 355
  iodide complexes, 157, 655
  iron(-II) carbonyl nitrosyl complex, 45
  iron(II) arene carbaborane complexes, 77
  iron(III) fluoride complexes, 737
  isocyanide complexes, 803
  ketoamine iron(III) polymer, 799
  lanthanide silylamide phosphine oxide complexes, 307
  lead(II) mercaptoethanol complex, 836
  macrocyclic diamine diimine complexes, 377
  magnesium amide dinitrogen and azide hydrazine complexes, 421
  mercaptopiperidine complexes, 157
  mercury(II) chloride complexes, 243
  mercury(II) mercapto complexes, 647
  mercury(II) organo compounds, 711
  metallated phosphine complex, 83
  methanol, 441
  methyl compounds, 397
  molybdenum(II) carbonyl Schiff base complexes, 233
IR spectra (cont)
  molybdenum(V) oxo complexes, 109, 163
  molybdenum(IV,IV,VI) cyanide oxo complex, 834
  morpholine complexes, 437
  nickel(II) crown ether complexes, 409
  nickel(II) macrocyclic complexes, 377
  nickel(II) molybdate, 331
  nickel(II) nitroacetate complexes, 247
  nitrate, 525
  nitrate complexes, 377, 437, 633
  nitric acid, 525
  nitroacetate complexes, 247
  nitrosyl complexes, 45, 209
  osmium cluster carbonyl hydrides, 105
  osmium(II) acyl complex, 83
  osmium(III) metallated phosphine coplex, 83
  osmium(IV) chloride complexes, 83
  osmium(VI) oxo complex, 83
  oxo complexes, 31, 83, 109, 163, 553
  oxoquinolinate complexes, 163
  palladium(II) halide complexes, 179, 543
  perchlorates, 539, 543, 633
  peroxide complexes, 21, 553
  phosphate esters, 683
  phosphine oxide complexes, 307, 355, 637
  phosphine selenide and phosphine sulphide complexes, 781
  phosphonium salts, 83
  phosphorus-nitrogen compounds, 535
  platinum carbonyl halide pyridine complexes, 689
  platinum polyfluoroamide, -amines and -imines, 129
  poly(iminoethylene)dithiocarbamate and its complexes, 225

  purine complexes, 143
  pyridine complexes, 689, 711
  rhenium phosphine complexes, 441
  rhenium amide and imide complexes, 31
  rhodanine complexes, 633
  rhodium(I) bis(diphenylphosphino)methane complexes, 809
  rhodium(I) carbonyl hydride complex, 570
  ruthenium(II) arene carbaborane complex, 77
  ruthenium(II) carbonyl complexes, 97
  ruthenium(II) difluorophosphate complex, 97
  Schiff base complexes, 233, 487
  silver(I) difluorophosphate complex, 97
  silylamide complexes, 307
  sulphur fluorine compounds, 723
  terpyridyl derivatives, synthetic intermediates, 303
  thiazole complexes, 557, 655
  thiocyanate complexes, 135, 377, 711
  thiourea complexes, 557
  tin(IV) organo halides, 397
  titanium(II) chloride complexes, 393
  trialkylammonium salts, 623
IR spectra (cont)
  tungsten (V) phenoxo complexes, 641
  uranium(III) cyclopentadienyls, 417
  uranium(VI) dioxo fluoride complex, 835
  vanadium nitrosyl complex, 209
  vanadium(0) carbonyl complex, 209
  water complex, 143
  ylide complexes, 539
  zirconium(IV) Schiff base complex, 487
Iridacyclobutanes, 773
Iridacyclobutene, 774
Iridacyclohexane, 771
Iridacyclopentanes, 741, 748, 757, 769, 771
Iridacyclopentenes, 745, 755, 774
Iridacyclopropenes, 757
Iridium(I), carbonylhydridotris(triphenylphosphine)-, 583
Iridium(III), *trans*-dichlorotetrapyridine- hydrogen dinitrate, 525
Iron carbonyl methylacrylate complex, formation of metallacycles, 751
Iron(-II), compound with iron - lead bond, 45
Iron(0)
  acetylnonacarbonyltri-, 842
  bis[1,2-triborane(7)]hexacarbonyldi-, 521
  cluster acyl alkoxide alkylidyne carbonyl hydride complexes, synthesis and X-ray crystal structures, 842
  ethylidynemethoxononacarbonyltri-, 842
  nonacarbonyldi-, insertion into carbon - carbon bonds, 742
  pentacarbonyl-, formation of metallocycles, 743
  pentacarbonyl-, reactiopn with tetracyanoethene, 799
  tricarbonyl[1-tetraborane(8)]-, 521
  tricarbonyl[2-tetraborane(8)]-, 521
Iron(II)
  amminopentacyano-, sodium salt trihydrate, 429
  aquopentacyano-, 429
  arene carbaborane complexes, synthesis, IR, mass and NMR(^{1}H, ^{11}B) spectra and X-ray crystal structure, 77
  bis(η^6-hexamethylbenzene)-, hexafluorophosphate, 77
  bisη^6-mesitylene-, hexafluorophosphate, 77
  bis(pentamethylcyclopentadienyl)-, 663
  bis[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
  chloride, hydrated complex with poly(iminoethylene)-dithiocarbamate, 225
  cyclopentadienyldicarbonyl(methyl-1-naphthylphenylstannyl)-, 1
  dimethylsulphoxidepentacyano-, 429
  dithiooxamidepentacyano-, 429
    hexamethylbenzene[-undecahydrodicarba-*nido*-undecaborate(2)], 77
  isonicotinamidepentacyano-, 429
  η^6-mesitylene-η^5-undecahydrodicarba-*nido*-undecaborate(2), 77
  organocompound, synthesis, IR, mass and NMR (^{1}H) spectra, 1
  oxidation by trismalonatocobaltate(III), kinetic study, 149
  pentacyano-, electrochemistry, electronic spectra, reaction with thioamides, kinetic study, 429

  pentacyano(thioacetamide)-, 429
Iron(II) (cont)
  pentacyano(thiourea)-, 429
Iron 3+, reaction with pyridine carboxylic acids, kinetic and thermodynamic study, 269
Iron(III)
  bis(2,6-dibenzothiazol-2-ylpyridine) chloride monohydrate, 543
  6,6''-bismethylhydrazino-4'-phenyl-2,2':6',2''-terpyridyl-, chloride tetrafluroborate monohydrate, 303
  chloride, hydrated complex with poly(iminoethylene)-dithiocarbamate, 225
  dichlorotetraaquo- dihydrate, 543
  difluoro(tetraphenylporphyrinato)-, tetrabutylammonium salt, 737
  fluoro porphyrin complexes, synthesis, electronic, ESR and IR spectra and magnetic properties, 737
  fluoro(tetraphenylporphyrinato)-, 737
  hydroxide ketoamine polymer, synthesis, IR and Moessbauer spectra and magnetic properties, 799
  pyridine-2-carboxylic acid complex, 269
  pyridine-2,6-dicarboxylic acid monoanion complex, 269
  tetrachloro-, methyltrioctylammonium and trioctylammonium salts, 623
  triaquotrifluoro-, 737
Isocyanide complexes, chromium(0), platinum(II), rhenium(I), tungsten(0) and (I), 803
Isoleucine complexes, uranium(VI), 537
Isolobal principle, boranes and ferraboranes, 521
Isomerism
  camphorquinonedioxime, 503
  chromium(III) β-diketonates, solvent extraction, 383
  copper(II) dioxime complex, 503
  gallium(III) and indium(III) β-diketonates, 9
  keto-enol, in tetradentate Schiff base, 487
  osmium gold cluster carbonyl hydride, 105
  palladium(II) thioether complexes, 289
  platinum(II) phosphine complexes, 89
  platinum(II) selenoether and thioether complexes, 289
  rhodium(I) and (III) vinyl complexes, 575
  rhodium(III) aminocarboxylate diamine complexes, 659
Isonicotinamide, reaction with iron(II) pentacyano thioamide complex, 429
Isotope exchange
  catalysis by rhodium(I) complex, 578
  hydrolysis of hydro(pyrrolyl)borates, 673
Isotopic substitution, germanium and tin organo halides, 397

K

Ketoamine complex, iron(III) polymer, 799
Kinetic study
  alkene hydrogenation catalysed by rhodium(I) complexes, 583
  alkene hydroformylation, 591, 593
  ammonolysis of trimetaphosphate, 425
  cobalt(III) base hydrolysis, 203
Kinetic study (cont)
  complex formation of iron(III) with pyridine carboxylic acids, 269
  copper(II) complexation by apotransferrin, 361
  copper(II) oxidation of iodide, 611
  copper(II) oxidation of trimethylphosphite, 467
  europium(III) aminopolycarboxylate complex reaction with ytterbium(3+), 707
  hydrolysis of fluoropolytungstates, 263
  hydrolysis of hydro(pyrrolyl)borates, 673
  inversion at pyramidal selenium and sulphur in palladium(II) and platinum(II) complexes, 289
  ligand exchange, manganese(II) prolinate complex, 113
  molybdenum(V) aquo oxo complexes, 109
  oxidation of iron(II) by cobalt(III) complex, 149
  oxidation of silver by ozone in alkali, 183
  platinum(II) substitution reactions, 201
  substitution reactions of aquopentacyanoferrate(II) with thioamides, 429

L

L-Lactate complex, terbium(III), 693
Lamellar compounds, zirconium(IV) sulphophosphonates, 61
Lanthanides
  chiral complexes, fluorescence and circularly polarised fluorescence spectra, Pfeiffer effect and formation constants, 195
  cyclopentadienyls, synthesis, 423
  diaminotetracarboxylate complexes, formation constants and use for cation exchange separation from actinides, 277
  dibutylphenacylphosphonate complexes, formation and use in separation by solvent extraction, 257
  fluoro hydroxo complexes, formation constants and geochemical significance, 239
  halide and nitrate complexes, formation constants, 49
  halides, oxides and hydrated ions, prediction of enthalpies of formation and disproportionation, 219
  silylamide phosphine oxide complexes, synthesis IR and NMR (^{1}H) spectra, 307
  toluene-4-sulphonate complexes, synthesis, NMR (^{1}H) spectra, electric conductance and solvation, 253
Lanthanide shifts (NMR)
  bistrimethylsilylamide, 307
  dimethylsulphoxide and toluene-4-sulphonate, 253
  trimethylphosphine oxide, 307
Lanthanum(III)
  bis(dibutylphenacylphosphonato)dibutylphenacylphosphonic acid perchlorato-, 257
  2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate complex, 277
  1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate complex, 277
  halide and nitrate complexes, formation constants, 49
  toluene-4-sulphonate, 253
  trimethylphosphine oxide tris(bistrimethylsilylamido)-, 307
Lead(II)
  bromide complexes in propylene carbonate, electronic spectra, electrochemistry and solubility, 169
  mercaptoethanol complexes, synthesis, IR spectra, formation constants and potentiometric titrations, 836
  mercaptoethanol-, 836
  mercaptoethanoldi-, 836
  pentakis(mercaptoethanol)tri-, 836
  tetrabromo-, 169
  tetrakis(mercaptoethanol)tri-, 836
  tribromo-, and undecabromotetra-, 169
Lead(IV)
  compounds with bonds to transition elements, synthesis and IR spectra, 45
  tetrakis(dicarbonylnitrosyltriphenylphosphiteiron)-, 45
  tetrakis(tricarbonyltri-*n*-butylphosphinecobalt)-, 45
  tetrakis(tricarbonyltriethylarsinecobalt)-, 45
  tetrakis(tricarbonyltriethylphosphinecobalt)-, 45
  tetrakis(tricarbonyltriphenylphosphite)-, 45
Lead(IV) (cont)
  tetrakis(tricarbonyltri-*n*-propylphosphinecobalt)-, 45
  tetrakis(tricarbonyltris{*n*-butyldiphenylphosphine}cobalt-, 45
  tetrakis(tricarbonyltris{diphenylmethylphosphine}cobalt-, 45
Leucine complex, uranium(VI), 537
Ligand exchange, manganese(II) prolinate complex, 113
Ligand reactivity, di-2-pyridyl ketone, cobalt(II) and (III) complexes, 21
Linalool, 594
Liquid-liquid partition coefficients, *see* solvent extraction
Lithium
  bromide, equilibria with lead(II) bromide, 169
  ditetrahydrofuran-, hexaphenoxytungstate(V), 641
  iodide, 629
  methyl, 441
  tetraphenylbuta-1,3-diene-1,4-di-, 549
Luminescence dissymmetry factors, terbium(III) complexes, 693
Lutetium(III)
  2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate complex, 277
  1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate complex, 277

M

Macrocyclic tetramine complexes, cobalt(II), copper(II), nickel(II), 377
Magnesacyclopentane, 748
Magnesium
  amide dinitrogen and azide hydrazine complexes, synthesis, electronic and IR spectra and thermogravimetric analysis, 421
  bromo(4-methoxyphenyl)-, 629
  diamidedinitrogen-, 421
  diazidodihydrazine-, 421
  polyphosphates, 69
Magnetic properties
  chromium(II) binuclear carboxylate, 135
  cobalt(II)
    macrocyclic tetramine complexes, 377
    nitroacetate complex, 247
    polypyridyl complexes, 21
    thiocyanate complexes, 711
  cobalt(III)
    binuclear complex, 21
    poly(iminoethylene)dithiocarbamate complex, 225
  copper(II)
    carboxylate complex, 679
    macrocyclic tetramine complexes, 377
    nitroacetate complexes, 247
    oxime complexes, 607
    poly(iminoethylene)dithiocarbamate complex, 225
    purine complexes, 143
  iron(II) poly(iminoethylene)dithiocarbamate complexes, 225
  iron(III)
    fluoride porphyrin complex, 737
  Magnetic properties [iron(III)] (cont)
    hydroxo ketoamine polymer, 799
    poly(iminoethylene)dithiocarbamate complexes, 225
  manganese(II)
    biacetylhydrazone complex, 831
    bipyridyl and bipyridyldioxide complexes, 327
    prolinate complex, 113
  nickel(II)
    macrocyclic tetramine complexes, 377
    nitroacetate complex, 247
    poly(iminoethylene)dithiocarbamate complex, 225
    thiocyanate complexes, 711
  niobium(0) and rhenium(0) complexes, 209
  rhenium(IV) imide complex, 37
  tantalum(0) and vanadium(0) complexes, 209
  vanadium(IV) poly(iminoethylene)dithiocarbamate complex, 225
L-Malate complex, terbium(III), 693
Maleic anhydride, Diels-Alder reaction with organoindium compound, 549
Malonate complex, cobalt(III), 149
L-Mandelate complex, terbium(III), 693
Manganese(I)
  cyclopentadienyltricarbonyl-, 663
  pentamethylcyclopentadienyltricarbonyl-, 663
  tricarbonyl[undecahydrodicarba-nido-undecaborate(2)]-, 663
Manganese(II)
  aminoacid complex, formation, magnetic properties, NMR (^{13}C) spectrum, relaxation, molecular structure and ligand exchange kinetics, 113
  biacetyldihydrazone complex, synthesis, electronic, ESR and IR spectra, electric conductance and magnetic properties, 831
  bipyridyl and bipyridyldioxide complexes, electronic, ESR and photoacoustic spectra and magnetic properties, 327
  4,4′-bipyridylbissulphato-, 327
  2,2′-bipyridyldichloro-, 327
  4,4′-bipyridyldichloro-, 327
  2,2′-bipyridyldiacetato-, 327
  4,4′-bipyridyldiacetato-, 327
  4,4′-bipyridyldioxidebissulphato-, 327
  2,2′-bipyridyldioxidedichloro-, 327
  4,4′-bipyridyldioxidedichloro-, 327

  - 2,2′-bipyridyldioxidedithiocyanato-, 327
  - 4,4′-bipyridyldioxidedithiocyanato-, 327
  - 2,2′-bipyridyldithiocyanato-, 327
  - 4,4′-bipyridyldithiocyanato-, 327
  - 2,2′-bipyridylsulphato-, 327
  - bis(biacetyldihydrazone)dichloro-, 831
  - 6,6′′-bismethylhydrazino-4′-phenyl-2,2′:6′,2′′- terpyridyl-, hexafluorophosphate monohydrate, 303
  - tris(L-prolinato)-, 113
  - zinc phosphate [$Mn_3Zn_9(PO_4)_8$], synthesis and neutron diffraction, 349
- Mass spectra
  - beryllium oxo nitrate, 820
  - boron subhalides, 134, 319
  - cyclopentadienyl complexes, 417
  - iron(II) arenecarbaborane complexes, 77
  - methylcyclopentadienyl complex, 417
  - nickel carbaborane complex, 833
  - organoiron compound, 1
  - organotellurium compounds, 13
  - organotin compounds, 1
  - osmium(VI) alkyl, 83
  - osmium - gold cluster carbonyl hydrides, 105
  - platinum carbonyl halide ylide complexes, 689
  - ruthenium(II) arenecarbaborane complexes, 77
  - sulphur alkyl fluoride, 723
  - terpyridyl derivatives and their synthetic intermediates, 303
  - tungsten(IV) alkoxy alkyl cyclopentadienyl complexes, 629
  - uranium(III) cyclopentadienyls, 417
  - zirconium(IV) tetradentate Schiff base complex, 487
- Mechanism
  - catalytic hydroformylation, 587
  - catalytic hydrogenation, 581
  - catalytic hydrosilylation, 600
  - *see also* kinetic study
- 2-Mercaptoethanol complexes, lead(II), 836
- Mercaptopiperidine complexes, mercury(II), 647
- Mercury, formation from di(*t*-butyl)mercury, 413
- Mercury(II)
  - alkyl, photochemical decomposition and EHMO calculation, 413
  - arsine carboxylate and arsine carboxylic acid halide complexes, synthesis, IR and NMR (^{1}H) spectra and thermogravimetric analysis, 667
  - aryl halide compounds, synthesis, IR and NMR (^{1}H) spectra and reactions, 243
  - aryl thiocyanate complexes with cobalt(II) and nickel(II), synthesis, electronic and IR spectra, electric conductance, magnetic properties and "softness", 711
  - 2,2′-bipyridylbis[diphenyl{pentacarbonylchromium(0)}-phosphido]-, 819
  - 2,2′-bipyridylbis[diphenyl{pentacarbonylmolybdenum(0)}-phosphido]-, 819
  - 2,2′-bipyridylbis[diphenyl{pentacarbonyltungsten(0)}-phosphido]-, 819
  - bis(2-dicyclohexylarsinobenzoato)-, ethanol solvate, 667
  - bis(2-dimethylarsinobenzoato)-, ethanol solvate, 667
  - bisdimethylsulphoxidebis[diphenyl{pentacarbonylchromium(0)}-phosphido]-, 819
  - bisdimethylsulphoxidebis[diphenyl{pentacarbonylmolybdenum(0)}-phosphido]-, 819
  - bisdimethylsulphoxidebis[diphenyl{pentacarbonyltungsten(0)}-phosphido]-, 819
  - bis(2-diphenylarsinobenzoato)-, ethanol solvate, 667
- Mercury(II) (cont)
  - bis[diphenyl{pentacarbonylchromium(0)}phosphido]-1,10-phenanthroline-, 819
  - bis[diphenyl{pentacarbonylmolybdenum(0)}phosphido]-1,10-phenanthroline-, 819
  - bis[diphenyl{pentacarbonyltungsten(0)}phosphido]-1,10-phenanthroline-, 819
  - bis[2-di(4′-tolyl)arsinobenzoato]-, ethanol solvate, 667
  - bis-μ-(1-methyl-4-mercaptopiperidine)tetrabromodi-, monohydrate, 157

  bis-μ-(1-methyl-4-mercaptopiperidine)tetraiododi-, monohydrate, 157
  bis(tri-4-tolylphosphine selenide)tetrabromodi-, 781
  bis(tri-4-tolylphosphine selenide)tetrachlorodi-, 781
  bis(tri-4-tolylphosphine selenide)tetraiododi-, 781
  bis(tri-4-tolylphosphine sulphide)dinitrato-, 781
  bromide, 543, 781
  bromotris(diphenylthiophosphinyl)methanido-, 137
  chloride, 543, 781
  chloro(2,5-dichlorophenyl)-, 243
  chloro-1,10-phenanthroline(2,3,4,6-tetrachlorophenyl)-, 243
  chloro-1,10-phenanthroline(2,3,5,6-tetrachlorophenyl)-, 243
  chloro-1,10-phenanthroline(2,4,6-trichlorophenyl)-, 243
  chloro(2,3,4,5-tetrachlorophenyl)-, 243
  chloro(2,3,4,6-tetrachlorophenyl)-, 243
  chloro(2,3,5,6-tetrachlorophenyl)-, 243
  chloro(2,3,4-trichlorophenyl)-, 243
  chloro(2,4,6-trichlorophenyl)-, 243
  chlorotris(diphenylthiophosphinyl)methanido-, 137
  2,6-dibenzothiazol-2-ylpyridinedibromo-, 543
  2,6-dibenzothiazol-2-ylpyridinedichloro-, 543
  dibromo(2-dicyclohexylarsinobenzoic acid)-, 667
  dibromo(2-diethylarsinobenzoic acid)-, 667
  di(*t*-butyl)-, 413
  dichloro(2-diethylarsinobenzoic acid)-, 667
  2,5-dichlorophenyl(methyl)-, 243
  2-dicyclohexylarsinobenzoic acid(diiodo)-, 667
  2-diethylarsinobenzoic acid(diiodo)-, 667
  di(1-methyl-2(2′-mercaptoethyl)piperidine)- perchlorate, 647
  di(1-methyl-3-mercaptomethylpiperidine)-, perchlorate, 647
  di(1-methyl-4-mercaptopiperidine)-, perchlorate, 647
  halide nitrate phosphine selenide and phosphine sulphide complexes, synthesis, electric conductance and IR spectra, 781
  halide phosphinylmethylide compounds, preparation and NMR (^{31}P) spectra, 137
  halide thiolate complexes, synthesis, IR spectra, electric conductance and X-ray diffraction, 157
  hexanitratotetrakis(triphenylphosphine selenide)tri-, 781
  iodide, 781
  iodotris(diphenylthiophosphinyl)methanido-, 137
Mercury(II) (cont)
  mercaptopiperidine complexes, synthesis, IR and Raman spectra and X-ray crystal structure, 647
  1-methyl-2(2′-mercaptoethyl)piperidine-, perchlorate hemihydrate, 647
  1-methyl-3-mercaptomethylpiperidine-, perchlorate monohydrate, 647
  1-methyl-4-mercaptopiperidine-, perchlorate, 647
  1-methyl-4-mercaptopiperidinetetrachlorodi-, 157
  methyl(2,3,4,5-tetrachlorophenyl)-, 243
  methyl(2,3,4,6-tetrachlorophenyl)-, 243
  methyl(2,3,5,6-tetrachlorophenyl)-, 243
  methyl(2,3,4-trichlorophenyl)-, 243
  methyl(2,4,6-trichlorophenyl)-, 243
  1-naphthyl(thiocyanato)-, 711
  perchlorate hexahydrate, 647
  1,10-phenanthrolinebis(2,3,4,5-tetrachlorophenyl)-, 243
  phosphido complexes, synthesis and NMR (^{31}P) spectra, 819
  thiocyanato(4-tolyl)-, 711
Mesityl(diphenylmethyl)chlorophosphine complex, platinum(II), 89
P-Mesityldiphenylmethylenephosphine complexes, platinum(II), 89
Mesitylene complexes, iron(II) and ruthenium(II), 77
Metallation reactions, 179, 441
Metalloboranes, cobalt and nickel, 133
Metallocarbaboranes, iron(II) and ruthenium(II), 77
Metal - metal bond
  chromium(II) - chromium(II), 135
  gold(I) - gold(I), 105
  gold(I) - osmium, 105
  molybdenum(II) - molybdenum(II), 209
  osmium - osmium, 105
  osmium - platinum(0), 105

  rhenium(I) - rhenium(V), 31
  rhodium(0) - rhodium(0), 586
  rhodium(I) - rhodium(I), 578
  rhodium(II) - rhodium(II), 637
  tungsten(I) - tungsten(I), 803
Metallacyclic compounds, review of synthesis and reactions, 739
Me$_4$-[16]-tetene complexes, cobalt(II), copper(II), nickel(II), 377
Methane, formation from ethanol, 839
Methanol
  photochemical decomposition, 839
  solvent extraction of chromium(III) β-diketonates, 383
  solvolysis of chromium(III) thiocyanate complex, 191
Methionine complexes, uranium(VI), 537
Methoxide complexes
  iron(0), 842
  tungsten(IV), 629
  tungsten(VI), 561
4-Methoxyacetophenone-2′-pyridylhydrazone complex, palladium(II), 179
2-Methoxybenzoate, organotellurium compound, 13
1-Methoxy-but-2-yne, 563
2-Methoxy-2,6-dimethyloct-7-ene, 594
4-Methoxyphenyl compound, tungsten(IV), 629
1-Methoxypropene, 563
3-Methoxypropene, 563
3-Methoxypropyne, 563
6-Methoxy-1-tetralone oxime complex, rhodium(I), 501
7-Methoxy-1-tetralone oxime complex, rhodium(I), 501
4-Methylacetophenone-2′-pyridylhydrazone complexes, palladium(II), 179
4-Methylanisole, 629
Methyl compounds
  boron cage, 134
  mercury(II), 243
  osmium(II) and (IV), 83
  phosphorus(V), 123
  platinum(II), 89, 289, 803
  rhenium(I), 441, 803
  rhenium(III), 441
  rhenium(V), 31
  rhodium(I), 809
  tellurium, 13
  tin, 1
  tin(IV), 209
  tungsten(IV), 629
  vanadium(II), 209
N-Methylcamphidine, 824
Methyl cyanide, *see* acetonitrile
Methylcyclopentadienide, potassium salt, 417
η^5-Methylcyclopentadienyl complex, uranium(III), 417
Methyldiphenylphosphine complexes
  cobalt(-I) and iron(-II), 45
  molybdenum(0), 130
4-Methylhexanal, 589
1-Methyl-2-(2′-mercaptoethyl)piperidine complexes, mercury(II), 647
1-Methyl-3-mercaptomethylpiperidine complexes, mercury(II), 647
1-Methyl-4-mercaptopiperidine complexes
  cadmium(II), 157
  mercury(II), 157, 647
  zinc(II), 157
Methyl methacrylate, polymerisation, 602
Methylnicotinate complex, chromium(III), 471
3-Methylocta-2,6-diene, hydroformylation, 593
3-Methylpent-2-ene, 589
5-Methylphenanthroline complex
  solvent extraction, 387
  zinc(II) complex, 387
4-Methylphenoxy compound, tungsten(IV), 629
4-Methylphenyl compound, tungsten(IV), 629
2-Methyl-2-phenylpropyl, tin compound, 1
1-[4′-Methylphenyl]-4,4,4-trifluoro-1,3-butanedionate, gallium(III) and indium(III) compounds, 9
2-Methylpropene, hydrogenation, 582

Methyl propiolate, formation of metallacycles, 753
N-Methylrhodanine complexes, copper(I) and silver(I), 633
1-Methylseleno-2-methylthioethane, platinum(II) complexes, 289
Methylthymol blue complexes, aluminium(III), 405
Methyltrimethoxyphosphonium ion, 467
Methyltrioctylamine, solvent extraction of metal chloride complexes, 623
Mixed metal cluster complexes
  gold(I) - osmium, 105
  osmium - platinum(0) 105
Mixed valence compounds
  molybdenum(IV, IV, VI), 834
  rhenium(I,V,V), 31
Moessbauer spectra
  iron(III) hydroxoketoamine polymer, 799
  tin(II) and (IV) dithiocarboxylate complexes, 187
  tin(II) organo compounds, 826
Molecular orbital theory, *see* EHMO
Molecular structure (not determined by diffraction methods), manganese(II) prolinate complex, 113
Molybdacyclobutanes, 770
Molybdate
  di-, sodium salt, 479
  β-octa-, ammonium salt pentahydrate, 17
Molybdenum, removal of rhenium and tungsten, 479
Molybdenum(0)
  bipyrimidine bridged binuclear complexes, synthesis, electronic, ESR, IR, and NMR (^{1}H, ^{13}C) spectra and redox reactions, 53
  μ-2,2′-bipyrimidine anion)octacarbonyldi-, 53
  (2,2′-bipyrimidine anion)tetracarbonyl-, 53
  μ-2,2′-bipyrimidine(octacarbonyl)chromium-, 53
  μ-2,2′-bipyrimidineoctacarbonyldi-, 53
  μ-2,2′-bipyrimidine(octacarbonyl)tungsten-, 53
  2,2′-bipyrimidinetetracarbonyl-, 53
  carbonyl phosphine complexes, NMR (^{95}Mo) spectra, 130
  carbonyl phosphine complexes, synthesis and NMR (^{31}P) spectra, 819
  pentacarbonyl(triphenylphosphine)-, 130
  tetracarbonylbis(methyldiphenylphosphine)-, *cis*- and *trans*-, 130
  tricarbonyltris(methyldiphenylphosphine)-, 130
  tris(2,2′-bipyridyl)-, 209
Molybdenum(II)
  carbonyl cyclopentadienyl pyrrole Schiff base complex, X-ray crystal structure and absolute configuration, 617
  η^5-cyclopentadienyldicarbonylpyridine-2-(2′-aza-4′,4′-dimethyl-1′,3′ -diphenylbut-1′-enyl)-, hexafluorophosphate salt, 233
  η^5-cyclopentadienyldicarbonylpyridine-2-(2′-aza-4′,4′-dimethyl-3′-phenylbut-1′-enyl)-, 233
  η^5-cyclopentadienyldicarbonylpyridine-2-(2′-aza-1′,3′-diphenyl-but-1′-enyl)-, hexafluorophosphate salt, 233
  η^5-cyclopentadienyldicarbonylpyridine-2-(2′-aza-1′,4′-diphenyl-3′-methyl ester but-1′-enyl)-, hexafluorophosphate salt, 233
  η^5-cyclopentadienyldicarbonylpyridine-2-(2′-aza-3′-methyl ester-pent-1′-enyl)-, 233
  η^5-cyclopentadienyldicarbonylpyridine-2-(2′-aza-3′-methyl ester-4′-phenylbut-1′-enyl)-, 233
Molybdenum(II) (cont)
  η^5-cyclopentadienyldicarbonylpyridine-2-(2′-aza-3′-methyl ester-1′-phenylpent-1′-enyl)-, hexafluorophosphate salt, 233
  η^5-cyclopentadienyldicarbonylpyridine-2-(2′-aza-1′-methyl-3′-methyl ester pent-1-enyl)-, hexafluorophosphate salt, 233
  η^5-cyclopentadienyldicarbonylpyridine-2-(2′-aza-1′-methyl-3′-methyl ester-4′-phenylbut-1-enyl)-, hexafluorophosphate salt, 233
  cyclopentadienyldicarbonylpyrrole-2-ald(2′-phenyl)ethylimine-, 617
  cyclopentadienyl(methyl)tricarbonyl-, 663
  methyl(pentamethylcyclopentadienyl)tricarbonyl-, 663
  methyltricarbonyl[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
  square pyramidal carbonyl cyclopentadienyl pyridine Schiff base complexes, synthesis, IR and NMR (^{1}H) spectra, conformational analysis and diastereoisomer equilibria, 233
  μ-tetraacetatodi-, 209

Molybdenum(IV)
  (2,2′-bipyridyl)tetrachloro-, synthesis, 209
  diacetonitriletetrachloro-, 209
  (1,10-phenanthroline)tetrachloro-, synthesis, 209
Molybdenum(IV,IV,VI)
  cyanide oxo complex, synthesis, electronic and IR spectra, 834
  hexaoxooctacyano-, potassium salt dihydrate, 834
Molybdenum(V)
  chloride, cocatalyst in carbonylation, 601
  di-μ-oxobis{(2-methyl-8-oxoquinoline)oxo-}, 163
  hydroxobis(2-methyl-8-oxoquinoline)oxo-, 163
  μ-oxobis{bis(2-methyl-8-oxoquinoline)oxo-}, 163
  oxo oxoquinoline complexes, synthesis, electronic and IR spectra, magnetic properties and DTA, 163
  polymeric aquo oxo ion, formation, sodium salt, electronic and IR spectra, kinetic study of conversion to dimer, 109
  tetraoxodi-, formation from polymeric ion, kinetic study, 109
Molybdenum(VI)
  dichlorodioxo-, 479
  nickel oxide (nickel molybdate), 331
  oxide, purification, reaction with sodium chloride, DTA and thermogravimetric study, 479
  oxotetrachloro-, 479
Morpholine, reaction with copper and hydrogen peroxide, 437
Morpholine complexes, copper(II), 437
Mulliken overlap populations, see overlap populations

N

^{14}N - ^{195}Pt coupling, in NMR (^{195}Pt) spectra of platinum(II) complexes, 57
Nalidixate complexes, copper(II), 153
1-Naphthyl compounds
  mercury(II), 711
  tin(IV), 1
1-[2′-naphthyl]-4,4,4-trifluoro-1,3-butanedionate, gallium(III) and indium(III) compounds, 9
Neodymium, reaction with thallium(I) cyclopentadienide, 423
Neodymium(III)
  complexes with 2,2′-diaminoether, N,N,N′,N′-tetraacetate and 1,5-diaminopentane,N,N,N′,N′-tetraacetate, 277
  halide and nitrate complexes, formation constants, 49
  tetrahydrofurantris(cyclopentadienyl)-, 423
  toluene-4-sulphonate, 253
Neopentyl complex, rhodium(I), 809
Neophyl complex, rhodium(I), 809
Nerolidol, 594
Neutron diffraction crystal structure
  copper(II) chloride complex, 365
  hexamminecobalt(III) salt, 365
Neutron powder diffraction
  aluminium(III) - titanium(II) chloride complex, 393
  manganese zinc phosphate, 349
Nickel
  carbaborane complexes, syntheses and mass and NMR (^{1}H, ^{11}B) spectra, 833
  η^5-cyclopentadienyl borane complex, bonding, 133
Nickel(I), 4′,4′′′-diphenyl-2,2′:6′,2′′:6′′,2′′′:6′′′,2′′′′-quinquepyridyl complex, 311
Nickel(II)
  bis(N-Benzothiazolyl-N′-ethylthiourea)-, 557
  bis(N-Benzothiazolyl-N′-methylthiourea)-, 557
  bis(2,2′-bipyridyl)nitroacetato-, trihydrate, 247
  bis(camphorquinone dioxime monoanion)-, 503
  biscyclopentadienyl-, 663, 833
  bis(2,6-dibenzothiazol-2ylpyridine)- perchlorate dihydrate, 543
  bisdimethylsulphoxidedi[1-naphthylmercury(II)thiocyanato]dithiocyanato-, 711
  bisdimethylsulphoxidedithiocyanatodi[4-tolylmercury(II)-thiocyanato]-, 711
  6,6′′-bismethylhydrazino-4′-phenyl-2,2′:6′,2′′- terpyridyl-, tetrafluoroborate hemihydrate, 303
  bis(pentamethylcyclopentadienyl)-, 663
  bis(1-phenyl-3-methyl-4-benzoylpyrazol-5-onato)-, 461

  bispyridinedi[1-naphthylmercury(II)thiocyanato]dithiocyanato-, 711
  bispyridinedithiocyanatodi[4-tolylmercury(II)thiocyanato]-, 711
  bis[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
  N,*N*′-butylenebis(2,4-pentanedioneiminato)-, 785
  N,*N*′-butylenebis(5,5,5-trifluoro-2,4-pentanedioneiminato)-, 785
Nickel(II) (cont)
  chloride, crown ether complexes, synthesis, IR spectra and X-ray crystal structure, 409
  chloride, hydrated complex with poly(iminoethylene)-dithiocarbamate, 225
  diacetatotetraaquo-, 557
  diaquodichlorodi-. 18-crown-6, 409
  dibromo(Me_4-[16]-tetene)-, 377
  dichloro(Me_4-[16]-tetene)-, 377
  di-μ-chlorooctaaquodi-, dichloride. 18-crown-6, 409
  dihistidine-, 791
  dihistidine(pyridoxamine)-, 791
  β-diketonate complexes, formation constants, solvent extraction and synergic effects, 461
  di[1-naphthylmercury(II)thiocyanato]dithiocyanato-, 711
  dinitrato(Me_4-[16]-tetene)-, 377
  di[nitroacetatobis(1,10-phenanthroline)-], heptahydrate, 247
  dioxime complex, formation, optical activity and solvent extraction, 503
  4′,4‴-diphenyl-2,2′:6′,2″:6″,2‴:6‴,2⁗-quinquepyridyl-, hexafluorophosphate, 311
  dipyridoxamine-, 791
  dipyridoxamine(histidine)-, 791
  dithiocyanatodi[4-tolylmercury(II) thiocyanato]-, 711
  dithiocyanato(Me_4-[16]-tetene)-, 377
  N,*N*′-ethylenebis(2,4-pentanedioneiminato)-, 785
  N,*N*′-ethylenebis(5,5,5-trifluoro-2,4-pentanedioneiminato)-, 785
  N,*N*′-ethylene[(2,4-pentanedioneimine)(5,5,5-trifluoro-2,4-pentanedioneiminato)]-, 785
  hexaquo-, chloride, 409
  hexaquo-, perchlorate, 543
  histidine-, 791
  histidine pyridoxamine complexes, electronic spectra, formation constants and potentiometric titrations, 791
  histidine(pyridoxamine)-, 791
  macrocyclic tetramine complexes, electronic and IR spectra, electric conductance and magnetic properties, 377
  molybdate, IR and Raman spectra, factor group analysis and structure, 331
  nitroacetate complexes, synthesis, electronic and IR spectra and magnetic properties, 247
  pyridoxamine-, 791
  N,*N*′-propylenebis(2,4-pentanedioneiminato)-, 785
  N,*N*′-propylenebis(5,5,5-trifluoro-2,4-pentanedioneiminato)-, 785
  tetradentate ketoimine complexes, synthesis, HPLC and isomerism, 785
  thiocyanate, complexes with mercury(II), synthesis, electronic and IR spectra, electric conductance, magnetic properties and "softness", 711
  tris(2,2′-bipyridyl)- dithiocyanato-1-naphthylmercurate(II), 711
  tris(2,2′-bipyridyl)- dithiocyanato-4-tolylmercurate(II), 711
  tris(1-phenyl-3-methyl-4-benzoylpyrazol-5-onato)-, trioctylammonium salt, 461
Nickel(IV)
  bis(pentamethylcyclopentadienyl)-, 663
  bis[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
Nickelacycloheptatriene, 759
Nickelacyclopentanes, 760, 770
Nickelacyclopentene, 767
Niobacyclopentadiene, 765
Niobacyclopentanes, 749, 765
Niobium(0)
  tris(2,2′-bipyridyl)-, magnetic properties, 209
  tris(1,10-phenanthroline)-, synthesis and magnetic properties, 209
Niobium(V), pentachloro-, 209
Nitrates and nitrate complexes
  beryllium, 820
  cobalt(II), 377

  copper(II), 377, 437
  lanthanum(III), 49
  mercury(II), 781
  neodymium(III), 49
  nickel(II), 377
  rhodium(III), 577
  silver(I), 633
Nitric acid (^{15}N), 525
Nitric acid, IR and Raman spectra, 525
Nitro compounds, carbonylation, 601
Nitroacetate complexes, cobalt(II), copper(II) and nickel(II), 247
4-Nitroacetophenone-2′-pyridylhydrazone complex, palladium(II), 179
Nitrobenzene, solvent for solvent extraction studies, 623
Nitrosyl complexes
  iron(-II), 45
  rhodium(III), 577
  vanadium, 209
7-Nitro-1-tetralone oxime complex, rhodium(I), 501
NMR (^1H) spectra
  acetoxyhydroborate complex, 299
  alkoxide complexes, 561
  alkyl complexes, 629
  amine complexes, 561
  aminopolycarboxylate complexes, 659
  aryl complexes, 243
  arylsalicylideneimine complexes, 213
  bipyrimidine complexes, 53
  bis(diphenylphosphino)methane complexes, 809
  cadmium(II) quinquepyridyl complexes, 311
  cobalamins and cobinamides, 321
  cobalt carbaborane complex, 833
  cobalt(III) binuclear complexes, 21
  cobalt(III) diaminoethane pentanedionate complex, 203
  copper(I) hydroborate phosphine complex, 299
  cyclopentadienyl complexes, 539, 629, 809
  cyclophosphamide complexes, 637
  diaminoethane complexes, 203, 659
  dimethylsulphoxide lanthanide complexes, 253
NMR (^1H) spectra (cont)
  ethanol, 667
  ethyl complex, 803
  hydride complexes, 441, 570, 629
  imido complexes, 561
  indium(III) organometallic compounds, 549
  iron(II) arene carbaborane complexes, 77
  isocyanide complexes, 803
  lanthanide silylamide phosphine oxide complexes, 307
  lanthanide toluene-4-sulphonate complexes, 253
  mercury(II) aryl complexes, 243
  metallated phosphine complex, 441
  methoxy compounds, 629
  methyl complexes, 629, 803
  methylcyclopentadienyl complex, 417
  molybdenum(II) cyclopentadienyl Schiff base complexes, 233
  neopentyl complex, 809
  neophyl complex, 809
  nickel carbaborane complex, 833
  nickel(II) quinquepyridyl complexes, 311
  organotellurium compounds, 13
  organotin compounds, 1
  osmium(II) acetyl complex, alkyls and phosphine complexes, 83
  osmium(III) alkyls, hydrido- and phosphine complexes, 83
  osmium(IV) alkyls and phosphine complexes, 83
  osmium(VI) alkyl, 83
  osmium(II,III) methylidene complex, 83
  palladium(II) aryl pyridylhydrazone complexes, 179
  palladium(II) thioether complexes, 289
  pentanedionate complex, 203
  phenyl compounds, 629
  phenylamido complex, 441
  phosphate esters, 683
  phosphine complexes, 31, 37, 97, 299, 441, 539, 803, 809, 819
  phosphine oxide complexes, 307

  phosphonium salts, 83
  platinum(II) ethylene complex, 57
  platinum(II) phosphine complexes, 89
  platinum(II) selenoether and thioether complexes, 289
  pyrazine complex, 417
  pyridine complexes, 803
  quinquepyridyl complexes, 311
  rhenium(I) and (III) amide and imide complexes, 37
  rhenium(I), (III) and (V) phosphine complexes, 441
  rhenium(V) amide and imide complexes, 31, 37
  rhodium(I) hydride complex, 570
  rhodium(I) oxime complexes, 501
  rhodium(III) aminocarboxylate complex, 659
  ruthenium(II) arene carbaborane complexes, 77
  Schiff base, 487
  selenoether platinum(II) complexes, 289
  silylamide complexes, 307
  terpyridyl derivatives, synthetic intermediates, 303
NMR (^{1}H) spectra (cont)
  tetra-n-butylammonium salt, 779
  1-tetralone oxime complexes, 501
  tetraphenylarsonium salts, 549
  thioether palladium(II) and platinum(II) complexes, 289
  tin(II) and (IV) cyclopentadienyldithiocarboxylate complexes, 187
  tin(IV) salicylideneimine complexes, 213
  toluene-4-sulphonate lanthanide complexes, 253
  trimethylphosphine complexes, 809
  trimethylsilylmethyl complex, 809
  triphenylphosphine complexes, 37, 539, 809, 819
  tungsten(V) phenoxo complexes, 641
  unsaturated diamine, 563
  uranium(III) cyclopentadienyls, 417
  ylide complexes, 539
  zirconium(IV) tetradentate Schiff base complex, 487
NMR (^{11}B) spectra
  cobalt carbaborane complex, 833
  copper(I) hydroborate complex, 299
  dicarbadecaborane(12) compounds, 822, 824
  iron(II) and ruthenium(II) arene carbaborane complexes, 77
  nickel carbaborane complexes, 833
NMR (^{13}C) spectra
  alkoxide complexes, 561
  alkylidyne complexes, 803
  amine complexes, 561
  arylsalicylideneimine complexes, 213
  bipyrimidine complex, 53
  carbon dioxide complex, 441
  carbonyl complex, 441
  cobalt(III) diaminoethane pentanedionate complex, 203
  cyclophosphamide complexes, 637
  diaminoethane complex, 203
  ethyl complex, 803
  gallium(III) and indium(III) β-diketonates, 9
  imido complexes, 561
  indium(III) organometallic complounds, 549
  isocyanide complexes, 803
  manganese(II) prolinate complex, 113
  metallated phosphine complex, 441
  methyl complex, 803
  organometallic compounds, book review, 139
  osmium(VI) alkyl, 83
  pentanedionate complex, 203
  phosphate esters, 683
  phosphine complexes, 441, 803
  silver(I) acetonitrile complexes, 145
  tin(IV) salicylideneimine complexes, 213
  trimethylphosphine complexes, 441
  unsaturated diamine, 563
NMR (^{19}F) spectra
  alumino-fluoropolytungstates, 263
  gallium(III) and indium(III) β-diketonates, 9
NMR (^{19}F) spectra (cont)
  polyfluoroamide, -amines and -imines, 129

  ruthenium(II) difluorophosphate and fluoride complexes and hexafluoroarsenate salt, 97
  trifluoromethyltin(II) compounds, 826
NMR (^{31}P) spectra
  amidotriphosphate, 425
  bis(diphenylphosphino)methane complexes, 809
  cyclophosphamide complexes, 637
  mercury(II) phosphine complexes, 819
  metallated phosphine complex, 441
  osmium(II) phosphine complexes, 83
  phosphate esters, 683
  phosphonium salts, 83, 123
  platinum(II) phosphine complexes, 89, 803
  polyphosphates, 69
  rhenium(I), (III) and (V) amide and imide complexes, 37
  rhenium(I), (III) and (V) phosphine complexes, 441
  rhenium(V) trimethylphospine complexes, 31
  ruthenium(II) difluorophosphate and phosphine complexes, 97
  silver(I) difluorophosphate and phosphine complexes, 97
  tetramethylphosphonium salt, 441
  trimethylphosphine complexes, 441, 803, 809
  tris(diphenylthiophosphinyl)methanido complexes, 137
NMR (^{75}As) spectrum, ruthenium(II) hexafluoroarsenate salt, 97
NMR (^{95}Mo) spectrum, molybdenum(0) carbonyl complexes, 130
NMR (^{109}Ag) spectrum, ruthenium(II)-silver(I) bonded complex, 97
NMR (^{195}Pt) spectra
  platinum(II) isocyanide methyl phosphine complex, 803
  platinum(II) and (IV) complexes, 57
NMR (NOE) spectrum, rhodium(I) bis(diphenylphosphino)methane triethylphosphine complex, 809
NMR (2-dimensional) spectra
  rhenium(I) metallated phosphine complex, 441
  rhodium(I) bis(diphenylphosphino)methane triethylphosphine complex, 809
Norbornadiene, hydroformylation, 592
Norbornadiene complexes, rhodium(I), 339
Normal coordinate analysis, germanium and tin organo halides, 397
Nuclear magnetic resonance, see NMR
Nuclear Overhauser effect, see NMR
Nucleophiles, reaction with platinum(II) complex, 201

O

Octabromooctaborane(8), 134
Octachlorononaborane(9), 134
Octachloro(methyl)nonaborane(9), 134
Octachlorooctaborane(8), 134
Octaiodooctaborane(8), 319
Octaphenyltetraphosphazene, 535
Octapolyphosphate, magnesium and tetramethylammonium salts, 69
Oct-1-ene, hydroformylation, 592
Optical activity
  camphorquinone dioxime, 503
  carbaborane, 824
  copper(II) and nickel(II) dioxime complexes, 503
  rhodium(III) aminocarboxylate diamine complexes, 659
Optical rotatory dispersion, rhodium(III) aminocarboxylate diamine complexes, 659
Orbital contour diagrams, boranes and ferraboranes, 521
Osmacyclopentadiene, 767
Osmacyclopentene, 744
Osmacyclopropene, 752
Osmium
  [bisdiphenylphosphinoethaneplatinum(0)]dodecacarbonyldihydridotetra-, 105
  carbohydrate polymers, preparation, density, viscosity, gel filtration and electrophoresis and sedimentation velocities, 335
  cluster carbonyl hydride with osmium – platinum bond, synthesis and IR spectrum, 105
  cluster carbonyl hydrides with gold phosphine moieties, synthesis, IR and mass spectra, X-ray crystal structure, isomerism and reaction, 105

  dioxo gluconate glucose polymers, 335
  dodecacarbonyldihydridobis[triethylphosphinegold(I)]tetra-, 105
  dodecacarbonyldihydridobis[triphenylphosphinegold(I)]tetra-,105
  dodecacarbonyl[triethylphosphinegold(I)]trihydridotetra-, 105
  dodecacarbonyltrihydrido[triphenylphosphinegold(I)]tetra-, 105
  dodecacarbonyltetrahydridotetra-, 105
Osmium(II)
  alkyls, synthesis, IR and NMR (^{1}H, ^{31}P) spectra and reactions, 83
  trans-bisacetyltetrakistriphenylphosphine-, 83
  trans-dichlorotetrakistrimethylphosphine-, 83
  trans-dimethyltetrakistriphenylphosphine-, 83
  phosphine complexes, synthesis, IR and NMR (^{1}H, ^{31}P) spectra and reactions, 83
Osmium(II,III)
  chloropentakistrimethylphosphinebis(μ-trimethylsilylmethylidene)di-, 83
  methylidene bridged binuclear complex, synthesis and NMR (^{1}H) spectra, 83
Osmium(III)
  alkyls, synthesis, ESR, IR and NMR (^{1}H) spectra, 83
  chloro(η^2-diphenylphosphinophenyl)hydrido(trimethylsilylmethyl)-triphenylphosphine-, 83
  η^2-diphenylphosphinophenyl(hydrido)methylbistriphenylphosphine-, 83
  η^2-diphenylphosphinophenyl(hydrido)(trimethylsilylmethyl)-triphenylphosphine-, 83
  phosphine complexes, synthesis, ESR, IR, and NMR (^{1}H) spectra and reactions, 83
  trichlorobistriphenylphosphine-, 83
Osmium(IV)
  alkyls, synthesis, IR and NMR (^{1}H) spectra, 83
Osmium(IV) (cont)
  chloride complexes, synthesis of phosphonium salts, IR and NMR (^{1}H, ^{31}P) spectra, electric conductance and reactions, 83
  hexachloro-, phosphonium salts, 83
  tetramethylbistriphenylphosphine-, 83
  trichlorobistrimethylphosphinetrimethylsilylmethyl-, 83
Osmium(VI)
  acetate oxo complex, X-ray crystal structure, 840
  alkyl, synthesis, IR, mass and NMR (^{1}H, ^{13}C) spectra, 83
  aqueous solutions, electronic spectra, 175
  dioxotriacetato-, potassium salt, 335
  formation in reduction of osmium(VIII) by hydrogen peroxide and in catalytic decomposition of hydrogen peroxide by osmium(VIII) oxide, 175
  oxotetrakistrimethylsilylmethyl-, 83
  triacetatodioxo-, potassium salt acetic acid solvate, 840
Osmium(VIII)
  aqueous solutions, electronic spectra, 175
  reduction by hydrogen peroxide, electronic spectra,
  electrochemistry and solvent extraction, 175
Overhauser effect, *see* NMR spectra
Overlap populations, boranes and ferraboranes, 521
Oxidative addition, rhodium(I) complex, 576
Oxide, molybdenum(VI)-nickel(II) ($NiMoO_4$), 331
Oxime, sulphonic acid complex, critical adsorption concentration, critical micelle concentration and interfacial tension, 453
Oxime complexes, rhodium(I), 501
Oxo complexes
  beryllium, 820
  molybdenum(IV,IV,VI), 834
  molybdenum(V), 109, 163
  osmium, 335
  osmium(VI), 83, 840
  rhenium(V), 31
  uranium(VI), 483, 537, 827, 835
2-Oxoacetophenoneoxime complex, copper(II), 607
2-Oxo-6-methylpyridinate complex, tungsten(I), 803
2-Oxo-1-naphthaldoxime complex, copper(II), 607
Oxoquinoline complexes, molybdenum(V), 163
1-Oxo-1,2,3,4-tetrahydronaphthalene oxime, *see* 1-tetralone oxime
Oxygen - bridged complex, rhenium(I,V), 31
Ozone, oxidation of silver in alkaline solutions, 183

P

^{31}P – ^{199}Hg coupling in NMR (^{31}P) spectra of mercury(II) complexes, 819
Palladacyclobutanes, 740, 774
Palladacyclopentadienes, 759
Palladacyclopentane, 747
Palladium, cocatalyst in hydroformylation, 592
Palladium(II)
  bisbenzonitriledichloro-, 543
  bis(pentamethylcyclopentadienyl)-, 663
  bis(triphenylphosphine)tetrachlorodi-, 539
  bis[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
  bromo[2-acetyl(2′-pyridylhydrazone)-5-chlorophenyl]-, 179
  bromo[2-acetyl(2′-pyridylhydrazone)-5-methylphenyl]-, 179
  bromo[3-acetyl(2′-pyridylhydrazone)-2-thiophenyl]-, 179
  N,*N*′-butylenebis(2,4-pentanedioneiminato)-, 785
  N,*N*′-butylenebis(5,5,5-trifluoro-2,4-pentanedioneiminato)-, 785
  chloro[2-acetyl(2′-pyridylhydrazone)-5-chlorophenyl]-, 179
  chloro[2-acetyl(2′-pyridylhydrazone)-5-methoxyphenyl]-, 179
  chloro[2-acetyl(2′-pyridylhydrazone)-5-methylphenyl]-, 179
  chloro[2-acetyl(2′-pyridylhydrazone)-5-nitrophenyl]-, 179
  chloro[2-acetyl(2′-pyridylhydrazone)-3-thiophenyl]-, 179
  chloro[3-acetyl(2′-pyridylhydrazone)-2-thiophenyl]-, 179
  chloro(cyclopentadienyl)triphenylphosphine-, 539
  cyclopalladation products from pyridylhydrazones of acetophenones and acetylthiophenes, synthesis, IR and NMR (^{1}H) spectra, 179
  cyclopentadiene phosphine ylide complexes, synthesis and IR and NMR (^{1}H) spectra, 539
  cyclopentadienyl(dimethylsulphonium benzoylmethylide)triphenylphosphine-, perchlorate, 539
  cyclopentadienyltriphenylphosphine(triphenylphosphonium acetylmethylide)-, perchlorate, 539
  cyclopentadienyltriphenylphosphine(triphenylphosphonium benzoylmethylide)-, perchlorate, 539
  2,6-dibenzothiazol-2-ylpyridinetetrachlorodi-, 543
  dibromo(3,4-dimethylthiotoluene)-, 289
  dichloro(1,2-dimethylthioethane)-, 289
  dichloro(1,2-dimethylthioethene)-, 289
  dichloro(3,4-dimethylthiotoluene)-, 289
  diiodo(1,2-dimethylthioethane)-, 289
  diiodo(1,2-dimethylthioethene)-, 289
  diiodo(3,4-dimethylthiotoluene)-, 289
  N,*N*′-ethylenebis(2,4-pentanedioneiminato)-, 785
  N,*N*′-ethylenebis(5,5,5-trifluoro-2,4-pentanedioneiminato)-, 785
  N,*N*′-ethylene[(2,4-pentanedioneiminato)(5,5,5-trifluoro-2,4-pentanedioneiminato)]-, 785
  iodo[2-acetyl(2′-pyridylhydrazone)-5-chlorophenyl]-, 179
  iodo[2-acetyl(2′-pyridylhydrazone)-5-methylphenyl]-, 179
  iodo[3-acetyl(2′-pyridylhydrazone)-2-thiophenyl]-, 179
  N,*N*′-propylenebis(2,4-pentanedioneiminato)-, 785
  N,*N*′-propylenebis(5,5,5-trifluoro-2,4-pentanedioneiminato)-, 785
Palladium(II) (cont)
  selenoether and thioether complexes, synthesis, NMR (^{1}H) spectra and fluxional behaviour, 289
  tetrabromo-, 179
  tetrachloro-, 179
  tetradentate ketoimine complexes, synthesis, HPLC and isomerism, 785
Papaverine complex, copper(II), 679
Partition coefficients, *see* solvent extraction
Penicillamine complex, cadmium(II), 153
Pentaborane(9), 521
Pentafluorophenyl complex, rhodium(I), 575
1-Pentafluorothiotetrafluoroethyl compound, sulphur(VI), 723
1- Pentafluorothio-2,2,2-trifluoroethyl compound, silver(I), 723
Pentamethylcyclopentadienyl, comparison with carbaborane, 663
neo-Pentane, formation in cyclometallation reactions, 772
2,4-Pentanedionate complexes
  cobalt(III), 203
  monodentate, 203
Pentapolyphosphate, magnesium and tetramethylammonium salts, 69
Pent-1-ene
  hydroformylation, 597

  isomerisation, 579
neo-Pentoxo complex, tungsten(VI), 561
Peonoloxime complex, copper(II), 607
Perbromate, caesium salt, 567
Perchlorate complex, silver(I), 97
Perfluoro- *see* heptafluoro-, hexafluoro- *etc.*
Peroxide, binuclear cobalt(III) complexes, 21
Pfeiffer effect, lanthanide complexes, 195
Phase transitions, tungsten(V) cyanide complex, 779
Phenanthridine di(nitric acid) dihydrate, 525
Phenanthrodinium hydrogen dinitrate and nitrate, 525
1,10-Phenanthroline
  monohydrate, 525
  reaction with mercury(II) aryl compounds, 243
1,10-Phenanthroline complexes
  chromium(0), 209
  cobalt(II) and copper(II), 247
  hafnium(0), 209
  mercury(II), 243, 819
  molybdenum(0) and (IV), 209
  nickel(II), 247
  niobium(0), rhenium(0), 209
  solvent extraction, 387
  tantalum(0), titanium(0), tungsten(0) and (IV), and vanadium(0), 209
  zinc(II), 387
  zirconium(0), 209
1,10-Phenanthrolinium
  hydrogen dinitrate, 525
  nitrate, 525
Phenoxy compound, tungsten(IV), 629
Phenoxymethyl compound, tungsten(IV), 629
Phenyl compounds
  palladium(II), 179
  phosphorus(V), 123
  tellurium, 13
  tin, 1
Phenylacetylene, 441
Phenylamide complexes
  rhenium(I) and (III), 37, 441
  rhenium(V), 31, 37
3-Phenylbut-1-ene, hydroformylation, 598
2-Phenyl-5,6-dicarba-*nido*-decaborane(12), 822
o-Phenylenebisdimethylarsine complex, molybdenum(0), 130
Phenylimide complexes
  rhenium(IV) and (V), 37
  tungsten(VI), 561
1-Phenyl-3-methyl-4-benzoylpyrazol-5-onate complexes, cobalt(II), indium(III), nickel(II), 457
1-Phenyl-3-methyl-4-lauroylpyrazol-5-onate complexes, indium(III), 457
1-Phenyl-3-methyl-4-(2-thenoyl)pyrazol-5-onate complexes, indium(III), 457
2-Phenylpropanal, 592, 598
1-Phenyl-4,4,4-trifluoro-1,3-butanedionate, gallium(III) and indium(III) compounds, 9
Phosphaalkene, platinum(II) complexes, 89
Phosphates and phosphate complexes, manganese(II), zinc(II), 349
Phosphate esters, synthesis, IR and NMR (^{1}H, ^{13}C, ^{31}P) spectra and uranium(VI) complexes, solvent extraction, 683
Phosphazenes, formation in reaction of chloramines with arylchlorophosphines, 535
Phosphines, reaction with chloramines, 535
Phosphonium salts
  bromochlorodiphenyl-, 123
  bromodichloro(phenyl)-, 123
  bromotrichloro-, 123
  bromotriphenyl-, tetrabromoborate, 123
  chlorotriethyl-, tetrachloroborate, 123
  chlorotrimethyl-, tetrachloroborate, 123
  chlorotriphenyl-, tetrachloroborate, 123
  dibromochloro(phenyl)-, 123

  - dibromodichloro-, 123
  - dibromodiphenyl-, tetrabromoborate, 123
  - dichlorodiethyl-, tetrachloroborate, 123
  - dichlorodimethyl-, hexachloroantimonate and tetrachloroborate, 123
  - dichlorodiphenyl-, tetrachloroborate, 123
  - hexachloroosmates(IV), synthesis, IR and NMR (^{1}H, ^{31}P) spectra, electric conductance and reactions, 83
  - synthesis and NMR (^{31}P) spectra, 123
  - tetrabromo-, 123
  - tetrabutyl-, chloride, 123
- Phosphonium salts (cont)
  - tetrachloro-, 123
  - tetraphenyl-, 123
  - tribromochloro-, 123
  - tribromo(phenyl)-, tetrabromoborate, 123
  - trichloro(ethyl)-, tetrachloroaluminate, 123
  - trichloro(methyl)-, tetrachloroaluminate, 123
  - trichloro(phenyl)-, 123
- Phosphorus(V)
  - alkyls, aryls and halocompounds, synthesis and NMR (^{31}P) spectra, 123
  - in heteropolytungstates, 283
- Photoacoustic spectra, manganese(II) bipyridyl and bipyridyl dioxide complexes, 327
- Photochemistry
  - mercury(II) alkyl, 413
  - rhodium(I) hydride phosphine complexes, 839
- 2-Phthalate, organotellurium compound, 13
- Picolinic acid, *see* pyridine-2-carboxylic acid
- Piperidine complexes, molybdenum(0), 130
- Plasma, *see* blood plasma
- Platinacyclobutenes, 739, 746, 772, 774
- Platinacycloheptane, 746
- Platinacycloheptatriene, 759
- Platinacyclohexanes, 746, 772
- Platinacyclopentanes, 746, 749, 768, 772
- Platinacyclopentenes, 745, 767, 772
- Platinacyclopropane, 746
- Platinum(0)
  - dihydridododecacarbonyltetraosmium(bisdiphenylphosphinoethane)-, 105
  - phosphine complex with osmium cluster carbonyl hydride, synthesis and IR spectrum, 105
  - tetrakistriphenylphosphine-, 592
- Platinum(II)
  - *cis*-bispyridinedichloro-, 525
  - *cis*-bis(trimethylphosphine)dimethyl-, 803
  - bromo(1,2-dimethylselenoethane)methyl-, 289
  - bromo(1,2-dimethylthioethane)methyl-, 289
  - bromo(methyl)(1-methylseleno-2-methylthioethane)-, 289
  - *t*-butylisocyanide-*cis*-dimethyl(trimethylphosphine)-, 803
  - *cis*-carbonyldibromo(1-pyridylpropylide)-, 689
  - *cis*-carbonyldichloro(1-pyridylpropylide)-, 689
  - carbonyl halide ylide complexes, synthesis,IR, mass and NMR (^{1}H) spectra and thermal decomposition, 689
  - *trans*-2-carboxypyridinedichloro(ethylene)-, 57
  - chloro(1,2-dimethylselenoethane)methyl-, 289
  - chloro(1,2-dimethylthioethane)methyl-, 289
  - chloro(methyl)(1-methylseleno-2-methylthioethane)-, 289
  - complexes, NMR (^{1}H, ^{195}Pt) spectra, 57
  - *cis*-diamminediaquo-, 57
  - *cis*-diamminedichloro-, 201
- Platinum(II) (cont)
  - (1,4-diazabicyclo-*trans*-undec-1-ene)dichloro(triethylphosphine)-, 89
  - dibromo(3,4-dimethylthiotoluene)-, 289
  - dibromo(pyridine)-1-pyridylpropylide-, 689
  - *trans*-dichloro(chloro⅝diphenylmethyl¾mesitylphosphine)triethylphosphine-, 89
  - *cis*-dichlorodi(*P*-mesityldiphenylmethylenephosphine)-, 89
  - dichloro(1,2-dimethylthioethane)-, 289

  dichloro(1,2-dimethylthioethene)-, 289
  dichloro(1,3-dimethylthiopropane)-, 289
  dichloro(3,4-dimethylthiotoluene)-, 289
  dichloro(*P*-mesityldiphenylmethylenephosphine)triethylphosphine-, *cis*- and *trans*- isomers and chloroform solvate of *cis*-isomer, 89
  dichloro(pyridine)-1-pyridylpropylide-, 689
  cis-diiododi(*P*-mesityldiphenylmethylenephosphine)-, 89
  diiodo(1,2-dimethylthioethane)-, 289
  diiodo(1,2-dimethylthioethene)-, 289
  diiodo(3,4-dimethylthiotoluene)-, 289
  cis-dimethyldi(*P*-mesityldiphenylmethylenephosphine)-, 89
  1,2-dimethylselenoethane(iodo)methyl-, 289
  1,2-dimethylthioethane(iodo)methyl-, 289
  iodo(methyl)(1-methylseleno-2-methylthioethane)-, 289
  isocyanide methyl phosphine complex, synthesis, IR and NMR (^{1}H, ^{13}C, ^{31}P) spectra, 803
  phosphaalkene complexes, synthesis, IR, NMR (^{1}H, ^{31}P) spectra and X-ray crystal structure, 89
  selenoether and thioether complexes, synthesis, NMR (^{1}H) spectra and fluxional behaviour, 289
  substitution reactions, kinetic study, 201
  tetraammine-, 57
  tetrapyridine-, nitrate, 525
  trichloro(triethylphosphine)-, hydro-1,5-diazabicyclo-[5,4,0]-undec-1-ene salt, 89
Platinum(IV)
  alkylcarbonyl halide ylide complexes, synthesis, IR and mass spectra, 689
  amine complex, NMR (^{195}Pt) spectrum, 57
  carbonylchloroiodo(methyl)pyridine(1-pyridylpropylide)-, chloride, 689
  cis, *trans*, *cis*-dichlorodihydroxobisisopropylamine-, 57
  dichloroiodo(methyl)pyridine(1-pyridylpropylide)-, 689
  trans-dichlorotetrapyridine-, nitrate, 525
  pyridine(1-pyridylpropylide)tetrachloro-, 689
Polarography
  aluminofluoroplytungstates and fluoropolytungstates, 263
  copper(II) morpholine complexes, 437
  heteropolytungstates, 283
  osmium(VI) and (VIII) in aqueous solutions, 175
Polyalkenes, formation of metallacycles, 765
Polyalkynes, formation of metallacycles, 765
Polyfluoroamide, -amines and -imines, synthesis, IR and NMR (^{19}F) spectra and reactions, 129
Poly(iminoethylene)dithiocarbamate and complexes, synthesis, electronic, ESR and IR spectra and magnetic properties, 225
Polyphosphates, protonation constants, NMR (^{31}P) spectra, anion exchange, magnesium and tetramethylammonium salts, 69
Polytungstates, 263
Porphyrin complexes, iron(III), 737
Potassium
  cyanide, hydroformylation catalyst, 596
  fluoride, 553
  methylcyclopentadienide, 1,2-dimethoxyethane solvate, 417
Potentiometric titrations
  aluminium(III) with methylthymol blue, 405
  cobalt(II) and copper(II) histidine pyridoxamine complexes, 791
  lead(II) bromide complexes, 167
  lead(II) mercaptoethanol complexes, 836
  nickel(II) histidine pyridoxamine complexes, 791
  osmium(VI) and (VIII) in aqueous solutions, 175
  thorium(IV) and uranium(VI) aminopolycarboxylate complexes, 827
  zinc(II) histidine pyridoxamine complexes, 791
Praseodymium(III)
  complexes with 2,2′-diaminoether*N*,*N*,*N*′,*N*′- tetraacetate and 1,5-diaminopentane*N*,*N*,*N*′,*N*′- tetraacetate, 277
  silylamide phosphine oxide complexes, 307
  toluene-4-sulphonate, 253
L-Proline complex, manganese(II), 113
Propane, formation from propene, 581
Propan-2-ol
  photochemical decomposition, 839

  solvolysis of chromium(III) thiocyanate complex, 191
Propan-2-ol-2-trimethylphosphonium, hexachloroosmate(IV), 83
Propene, 689
  hydroformylation, 585, 595
N-Prop-2-enylacetamide, hydroformylation, 599
Propionate complexes
  copper(II), 679
  rhodium(II), 637
i-Propoxide complexes, tungsten(VI), 561
2-Propylamine complex, platinum(IV), 57
N,*N*′-Propylenebis(2,4-pentanedioneimine) complexes, copper(II), nickel(II), palladium(II), 785
N,*N*′-Propylenebis(5,5,5-trifluoro-2,4-pentanedioneimine) complexes, copper(II), nickel(II), palladium(II), 785
Propylenecarbonate
  solvent extraction of chromium(III) complexes, 383
  solvent for lead(II) bromide complexes, 169
Propylenediaminetetraacetate complexes, europium(III) and ytterbium(III), 707
N,*N*′-Propylene[(2,4-pentanedioneimine)(5,5,5-trifluoro-2,4-pentanedioneimine)] complexes, copper(II), nickel(II), palladium(II), 785
4-Propyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane complexes, rhodium(I), 572
Protonation constants
  1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate, 277
  polyphosphates, 69
^{195}Pt - ^{31}P coupling, in NMR (^{31}P) spectra of platinum(II) complexes, 89
Purine complexes, copper(II), 143, 153
Pyrazine complexes
  bridged, 417
  uranium(III), 417
Pyridine complexes
  boron, 673
  cobalt(II), 21, 711
  cobalt(III), 21, 525
  iridium(III), 525, 549
  nickel(II), 711
  platinum(II) and (IV), 525, 689
  rhodium(I), 339
  rhodium(III), 525
  ruthenium(II) and (III), 525
Pyridine
  2-(2′-aza-4′,4′-dimethyl-1′,3′-diphenylbut-1′-enyl)-, molybdenum(II) complex, 233
  2-(2′-aza-4′,4′-dimethyl-3′-phenylbut-1′-enyl)-, molybdenum(II) complex, 233
  2-(2′-aza-1′,3′-diphenylbut-1′-enyl), molybdenum(II) complex, 233
  2-(2′-aza-1′,4′-diphenyl-3′-methyl ester but-1′-enyl)-, molybdenum(II) complex, 233
  2-(2′-aza-3′-methyl ester pent-1′-enyl)-, molybdenum(II) complex, 233
  2-(2′-aza-3′-methyl ester-4′-phenylbut-1′-enyl)-, molybdenum(II) complex, 233
  2-(2′-aza-3′-methyl ester-1′-phenylpent-1′-enyl)-, molybdenum(II) complex, 233
  2-(2′-aza-1′-methyl-3′-methyl ester pent-1′-enyl)-, molybdenum(II) complex, 233
  2-(2′-aza-1′-methyl-3′-methyl ester-4′-phenylbut-1′-enyl)-, molybdenum(II) complex, 233
Pyridine-2-carboxylic acid complex, iron(III), 269
Pyridine-2,6-dicarboxylate complexes, europium(III), holmium(III) and terbium(III), 195
Pyridine-2,6-dicarboxylic acid monoanion complex, iron(III), 269
Pyridoxamine complexes, cobalt(II), copper(II), nickel(II), zinc(II), 791
2-Pyridylhydrazone complexes, palladium(II), 179
Pyrogallol dianion complexes, aluminium(III), 313
Pyrophosphate, magnesium and tetramethylammonium salts, 69
Pyrrole complex, molybdenum(II), 617

Q

Quinoline complexes
  copper(II), 679

  rhodium(I), 339
Quinquepyridyl complexes, synthesis, ESR and NMR (^{1}H) spectra and electrochemistry, 311

R

Radiochemical study, zinc complexes, solvent extraction, 387
Raman spectra
  chloride complexes, 355, 475, 525
  chromium(III) chloride, 475
  fluoroalkyl compounds, 723
  fluorothio compounds, 723
  hexamethylphosphoramide complex, 355
  hydrogen dinitrate, 525
  indium(III) chloride phosphine oxide complex, 355
  mercury(II) mercaptide complexes, 647
  nickel molybdate, 331
  nitrate, 525
  phosphine oxide complex, 355
  pyridine complexes, 525
  sulphur fluorine compounds, 723
  uranium apatites, 343
  uranium(VI) dioxo fluoride complex, 835
Rate study, *see* kinetic study
Relaxation, NMR (^{13}C) spectrum of manganese(II) prolinate complex, 113
Resacetophenoneoxime complex, copper(II), 607
Review
  metallacyclic compounds, 739
  rhodium(I) carbonyl hydride phosphine complex, 569
Rhenium, removal from molybdenum, 479
Rhenium(0)
  tris(2,2′-bipyridyl)-, synthesis and magnetic properties, 209
  tris(1,10-phenanthroline)-, synthesis and magnetic properties, 209
Rhenium(I)
  bis(*t*-butylisocyanide)bis(trimethylphosphine)-*t*-butylaminomethylidyne(chloro)-, tetrafluoroborate, 803
  trans-bis(*t*-butylisocyanide)chlorotris(trimethylphosphine)-, 803
  biscarbonyl(chloro)tris(trimethylphosphine)-, 441
  μ-1,2-bis(η^5-3,4-diphenylcyclopentadienediyl)-1,2-diphenylethanedi[tris(trimethylphosphine)-], 441
  trans-bis(trimethylphosphine)chlorotris(*t*-butylisocyanide)-, 803
  t-butylaminomethylidyne(*t*-butylisocyanide)chlorotris(trimethylphosphine)-, tetrafluoroborate, 803
  η^2-carbon dioxide(phenylamido)tris(trimethylphosphine)-, 441
  chloro(dinitrogen)tetrakis[diethyl(phenyl)phosphine]-, 441
  chloropentakis(*t*-butylisocyanide)-, 803
  chloropentakis(trimethylphosphine)-, 441
  cyclopentadienyltricarbonyl-, 663
  dicarbonyl(phenylamido)tris(trimethylphosphine)-, 37
  η^2-dimethylphosphinomethyltetrakis(trimethylphosphine)-, 441
  dinitrogen(hydrido)tetrakis[diethyl(phenyl)phosphine]-, 441
  dinitrogenpentakis(trimethylphosphine)-, chloride methanol solvate, 441
  dinitrogenpentakis(trimethylphosphine)-, tetrafluoroborate, 441
  dinitrogen(phenylamido)bis(trimethylphosphine)-, 37
  dinitrogen(phenylamido)tetrakis(trimethylphosphine)-, 441
Rhenium(I) (cont)
  ethylpentakis(*t*-butylisocyanide)-, 803
  hydridopentakis(trimethylphosphine)-, 441
  isocyanide complexes, synthesis, IR and NMR (^{1}H, ^{13}C, ^{31}P) spectra, electric conductance and X-ray crystal structure, 803
  methylpentakis(*t*-butylisocyanide)-, 803
  methylpentakis(trimethylphosphine)-, 441
  μ-oxo-oxobis(trimethylsilylmethyl)rhenium(V)tetrakis(trimethylphosphine)[dioxo(trimethylsilylmethyl)rhenium(V)(Re-Re)]-, tetrahydrofuran solvate, 31
  pentakis(trimethylphosphine)-, chloride, 441
  pentamethylcyclopentadienyltricarbonyl-, 663
  phenylamido complexes, synthesis, IR and NMR (^{1}H, ^{31}P) spectra, X-ray crystal structure and reactions, 37
  phosphine complexes, synthesis, IR and NMR (^{1}H, ^{13}C, ^{31}P, 2-D)

    spectra, electric conductance and X-ray crystal structure, 441
  tricarbonyl(phenylamido)bis(trimethylphosphine)-, 37
  tricarbonyl[undecahydrodicarba-nido-undecaborate(2)]-, 663
Rhenium(III)
  chloro(hydrido)pentakis(trimethylphosphine)-, tetrafluoroborate, 441
  dihydrido(phenylamido)tetrakis(trimethylphosphine)-, 37
  η^1-dimethylphosphidomethyl(hydrido)phenylamidotetrakis(trimethylphosphine)-, 37
  iodo(methyl)tetrakis(trimethylphosphine)-, iodide, tetramethylphosphonium iodide, 441
  iodo(phenylamido)tetrakis(trimethylphosphine)-, 441
  phenylamido complexes, synthesis, IR and NMR (^{1}H, ^{31}P) spectra, 37
  phosphine complexes, synthesis, IR and NMR (^{1}H, ^{31}P) spectra, electric conductance and X-ray crystal structure, 441
  tetrakis(trimethylphosphine)trihydrido-, 441
Rhenium(IV)
  bistetrahydrofurantetrachloro-, 209, 441, 803
  dichloro(phenylamido)tristrimethylphosphine-, 37
  phenylimide complex, synthesis, ESR and IR spectra and magnetic properties, 37
Rhenium(V)
  alkyl compounds, synthesis, IR and NMR (^{1}H, ^{31}P) spectra, electric conductance and reactions, 31
  η^4-buta-1,3-diene(phenylamido)tris(trimethylphosphine)-, 37
  chloro(dimethyl)phenylimidobis(trimethylphosphine)-, 31
  diacetato(methyl)(phenylimido)bis(trimethylphosphine)-, 31
  dichloro(methyl)phenylimidobis(trimethylphosphine)-, 31
  dimethyl(phenylimido)bis(trimethylphosphine)-, tetrafluoroborate, 31
  fluoro(dimethyl)phenylimidobis(trimethylphosphine)-, tetrafluoroborate, 31
  oxobis(trimethylsilylmethyl)[μ-oxotetrakis(trimethylphosphine)-rhenium(I)]dioxotrimethylsilylmethylrhenium(V)(Re-Re)-, tetrahydrofuran solvate, 31
  phenylamide complex, synthesis, IR and NMR (^{1}H, ^{31}P) spectra and X-ray crystal structure, 37
Rhenium(V) (cont)
  phenylimide compounds, synthesis, IR and NMR (^{1}H, ^{31}P) spectra, electric conductance and reactions, 31, 37
  phenylimido(trimethyl)bis(trimethylphosphine)-, 31
  phosphine complex, synthesis, IR and NMR (^{1}H, ^{31}P) spectra and electric conductance, 441
  tetrahydridotetrakis(trimethylphosphine)-, tetrafluoroborate, 441
  trichloro(phenylimido)bistrimethylphosphine-, 37
  trichloro(phenylimido)trimethylphosphine(triphenylphosphine)-, toluene solvate, 37
Rhenium(I,V,V), trinuclear complex, synthesis, IR and NMR(^{1}H, ^{31}P) spectra and X-ray crystal structure, 31
Rhodacyclobutanes, 742, 754
Rhodacycloheptane, 748
Rhodacyclohexanes, 741, 748
Rhodacyclopentadienes, 754, 767
Rhodacyclopentanes, 740, 748, 755, 757, 769
Rhodacyclopentenes, 745, 754
Rhodanine complexes, copper(I) and silver(I), 633
Rhodium(0)
  dicarbonyltetrakis(triphenylphosphine)di-, 586
  tetracarbonyltetrakis(triphenylphosphine)di-, 585
Rhodium(I)
  alkene ,phosphine pyridine complexes, synthesis and electric electronic conductance, 339
  arene complex, formation from metallacycle, 754
  bis[bis(diphenylphosphino)methane]chloro-, 809
  bis(diphenylphosphino)methanechloro(trimethylphosphine)-, 809
  bis(diphenylphosphino)methanechloro(triphenylphosphine)-, 809
  bis(diphenylphosphino)methane complexes, synthesis, IR and NMR (^{1}H, ^{31}P, 2-D and NOE) spectra and X-ray crystal structure, 809
  bis(diphenylphosphino)methane(cyclopentadienyl)-, 809
  bis(diphenylphosphino)methane(cyclopentadienyl)-trimethylphosphine-, 809
  bis(diphenylphosphino)methane(methylcyclopentadienyl)-, 809

  bis(diphenylphosphino)methane(neopentyl)trimethylphosphine-, 809
  bis(diphenylphosphino)methane(neophyl)trimethylphosphine-, 809
  bis(diphenylphosphino)methane(trimethylphosphine)-trimethylsilylmethyl-, 809
  bis(2-ethylpyridine)1,5-cyclooctadiene-, tetraphenylboronate, 339
  bispyridine-1,5-cyclooctadiene-, tetraphenylboronate, 339
  bisquinoline-1,5-cyclooctadiene-, tetraphenylboronate, 339
  bisquinoline(norbornadiene)triphenylphosphine-, tetraphenylboronate, 339
  bisquinoline(norbornadiene)tri-4-fluorophenylphosphine-, tetraphenylboronate, 339
  bisquinoline(norbornadiene)tri-4-tolylphosphine-, tetraphenylboronate, 339
  bis(triphenylphosphine)bromo(carbonyl)-, 596
  bis(triphenylphosphine)carbonyl(chloro)-, 569, 593
  bis(triphenylphosphine)carbonyl(hydrido)-, 571
Rhodium(I) (cont)
  bis(triphenylphosphine)dicarbonyl(hydrido)-, 587
  carbonyl chloride oxime complexes, synthesis, NMR (^{1}H) spectra and carbon-hydrogen bond activation, 501
  carbonyl(hydrido)tris[diphenyl(trimethylsilylmethyl)phosphine]-, 591
  carbonyl(hydrido)tris(triphenylphosphine), review, 569
  carbonyltris(tri-*i*-propylphosphine)-, 839
  catalysis of quadricyclene isomerisation, 741
  chlorodicarbonyl(6-methoxy-1-tetralone oxime)-, 501
  chlorodicarbonyl(7-methoxy-1-tetralone oxime)-, 501
  chlorodicarbonyl(7-nitro-1-tetralone oxime)-, 501
  chlorodicarbonyl(1-tetralone oxime)-, 501
  chlorotetracarbonyldi-, insertion into carbon-carbon bonds, 740
  chlorotetrakis(trimethylphosphine)-, 809
  chlorotris(triphenylphosphine)-, 583, 809
  1,5-cyclooctadiene(2-ethylpyridine)triphenylarsine-, tetraphenylboronate, 339
  1,5-cyclooctadiene(2-ethylpyridine)triphenylphosphine-, tetraphenylboronate, 339
  1,5-cyclooctadiene(pyridine)triphenylphosphine-, tetraphenylboronate, 339
  1,5-cyclooctadiene(quinoline)triphenylarsine-, tetraphenylboronate, 339
  1,5-cyclooctadiene(quinoline)triphenylphosphine-tetraphenylboronate, 339
  dicarbonyl(hydrido)triphenylphosphine-, 588
  dichloro(dinorbornadiene)di-, 339
  2-ethylpyridine(norbornadiene)triphenylphosphine-, tetraphenylboronate, 339
  hydride phosphine complexes, photolysis in alcohols, formation of alkanes, 839
  hydridotris(triphenylphosphine)-, 570
  hydridotris(tri-*i*-propylphosphine)-, 839
  norbornadiene(pyridine)triphenylphosphine-, tetraphenylboronate, 339
  norbornadiene(quinoline)tri-3-tolylphosphine-, tetraphenylboronate, 339
  pentacarbonyltetra(μ-diphenylphosphido)triphenylphosphinetetra-, 587
Rhodium(II)
  carboxylate cyclophosphamide complexes, synthesis, electronic, IR and NMR (^{1}H, ^{13}C, ^{31}P) spectra, 637
  di(cyclophosphamide)tetraacetatodi-, 637
  di(cyclophosphamide)tetrabutyratodi-, 637
  di(cyclophosphamide)tetrapropionatodi-, 637
Rhodium(III)
  aminopolycarboxylate diamine complexes, synthesis, circular dichroism, electronic, NMR (^{1}H) and optical rotatory dispersion spectra and isomerism, 659
  cis-bis(bipyridyl)dichloro-, hydrogen trinitrate, 525
  1,2-diaminoethane(*N*,*N*′-dipropionato-1,2-diaminoethane)-, chloride, and monohydrate, 659
Rhodium(III) (cont)
  2,6-dibenzothiazol-2-ylpyridinetrichloro-, dihydrate, 543
  trans-dichlorotetrapyridine-, chloride pentahydrate and hydrogen dinitrate, 525

  metallacyclic compounds, 740
  triaquotrichloro-, 543
Rubber polymers, hydrogenation, 584
Ruthenacyclobutane, 772
Ruthenacyclobutene, 752
Ruthenacyclohexadiene, 753
Ruthenacyclopentadienes, 752
Ruthenacyclopentane, 754
Ruthenium(II)
  arenecarbaborane complex, synthesis, IR, mass and NMR (^{1}H, ^{11}B) spectra, 77
  benzene[undecahydrodicarba-*nido*-undecaborate(2)]-, 77
  trans-bis(bisdiphenylphosphinoethane)carbonyl(fluoro)-, 97
  trans-bis(bisdiphenylphosphinoethane)dicarbonyl-, hexafluoroantimonate, 97
  trans-bis(bisdiphenylphosphinoethane)dicarbonyl-, tetrafluoroborate, 97
  trans-bis(bisdiphenylphosphinomethane)carbonyl-(difluorophosphato)-, hexafluorophosphate, 97
  cis- and *trans*-bis(bisdiphenylphosphinomethane)dicarbonyl-, hexafluoroantimonate, 97
  cis-bis(bisdiphenylphosphinomethane)dicarbonyl-, hexafluorophosphate, 97
  cis-bis(bisdiphenylphosphinomethane)dicarbonyl-, tetrafluoroborate, 97
  cis-bis(bisdiphenylphosphinomethane)dichloro-, improved synthesis, 97
  cis-bisdiphenylphosphinomethanedicarbonyl-μ-bisdiphenylphosphinomethane[chlorosilver(I)]-, hexafluoroarsenate, 97
  cis-bisdiphenylphosphinomethanedicarbonyl-μ-bisdiphenylphosphinomethane[(difluorophosphato)- silver(I)]-, difluorophosphate 97
  cis-bisdiphenylphosphinomethanedicarbonyl-μ-bisdiphenylphosphinomethane[(fluorotrihydroxyborato)silver(I)]-, tetrafluoroborate, 97
  cis-bisdiphenylphosphinomethanedicarbonyl-μ-bisdiphenylphosphinomethane[perchloratosilver(I)]-, perchlorate, 97
  complexes with ruthenium - silver bonds, synthesis, IR and NMR (^{19}F, ^{31}P) spectra and electric conductance, 97
  dibenzenetetrachlorodi-, 77
  dicationic carbonyl phosphine complexes, synthesis,IR and NMR (^{19}F, ^{31}P) spectra and electric conductance, 97
  trans-dichlorotetrapyridine-, monohydrate, 525
  μ-difluorophosphatobis[*trans*-bis(bisdiphenylphosphinomethane-], hexafluorophosphate salt, 97
Ruthenium(III)
  2,6-dibenzothiazol-2-ylpyridine(trichloro)-, ethanol solvate, 543
  trans-dichlorotetrapyridine-, hydrogen dinitrate, 525
  trichloride trihydrate, 543

S

Safety
  boron subhalide explosion, 134
  perchlorate complex explosion, 97
Salicyl complex, zirconium(IV), 487
Salicylaldehyde, 487
Salicylate complexes, terbium(III), 693
N-Salicylideneaniline complex, tin(IV), 213
N-Salicylideneanisidine complex, tin(IV), 213
Samarium, reaction with thallium(I) cyclopentadienide, 423
Samarium(II) fluoride, improbability of disproportionation, 219
Samarium(III)
  2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate complex, 277
  1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate complex, 277
  tetrahydrofurantriscyclopentadienyl-, 423
  toluene-4-sulphonate, 253
Scandium(III)
  fluoride, reaction with hydrazine, IR spectra, 735
  hydrazinetrifluoro-, 735
  hydroxotrifluoro-, hydrazinium salt, 735
Scanning electron microscopy, *see* electron microscopy
Schiff base complexes
  copper(II), 785

  molybdenum(II), 617
  nickel(II) and palladium(II), 785
  zirconium(IV), 487
Sedimentation velocities, osmium carbohydrate polymers, 335
Selective hydrogenation catalysed by rhodium(I) complex, 581
Selenium, pyramidal, inversion barriers, 289
Selenoether complexes, platinum(II), 289
Semibullvalenes, insertion reactions, 742
Serine complexes, uranium(VI), 537
Silanes, in hydrosilylations, 600
Silica, hydroformylation catalyst support, 596
Silicon in heteropolytungstates, 283
Silicostannotungstate$(HSiSnW_{11}O_{40})^{5-}$, potassium salt hydrate, 283
Silver, oxidation by ozone in alkali, kinetic and electron microscopy studies, 183
Silver(I)
  acetonitrile-, 145
  acetonitrile complexes, NMR (^{13}C) spectra and formation constants, 145
  N-aminorhodaninenitrato-, 633
  bisacetonitrile-, 145
  complexes with ruthenium - silver bonds, synthesis, IR and NMR (^{19}F, ^{31}P) spectra and electric conductance, 97
  di(*N*-aminorhodanine)-, perchlorate, 633
  2,6-dibenzothiazol-2-ylpyridine-, perchlorate, 543
  cis-dicarbonylbisdiphenylphosphinomethane-μ-bisdiphenylphosphinomethaneruthenium(II)(difluorophosphato)-, difluorophosphate, 97
Silver(I) (cont)
  cis-dicarbonylbisdiphenylphosphinomethane-μ-bisdiphenylphosphinomethaneruthenium(II)(fluorotrihydroxyborato)-, tetrafluoroborate, 97
  cis-dicarbonylbisdiphenylphosphinomethane-μ-bisdiphenylphosphinomethaneruthenium(II)(perchlorato)-, perchlorate, 97
  N-methylrhodaninenitrato-, 633
  oxide, intermediate in oxidation of silver by ozone in alkali, 183
  1-pentafluorothiotetrafluoroethyl-, acetonitrile solvate, 723
  1-pentafluorothio-2,2,2-trifluoroethyl-, acetonitrile hemisolvate, 723
  perchlorate, 539, 543
  rhodanine complexes, synthesis, electric conductance and IR spectra, 633
Silver(II) oxide, formation in oxidation of silver by ozone, electron microscopy, 183
Sodium
  acetoxytrihydroborate, IR spectrum and reaction with copper(I) complex, 299
  amalgam, reduction of rhenium compounds, 441
  amalgam, synthesis of zerovalent compounds, 209
  chloride, 479
  cyclopentadienide, 1,2-dimethoxyethane solvate, 417
  fluoride, 553
  hydrotri*iso*propoxyborate, 299
" Softness", thiocyanate complexes, 711
Solubility study, lead(II) bromide complexes, 169
Solvent extraction
  chloride complexes, 623
  chromium(III) β-diketonates, isomers, 383
  cobalt(II) β-diketonate complexes, 461
  copper(II) with camphorquinone dioxime, 503
  heterocyclic diamines, synergic effect, 387
  indium(III) β-diketonate complexes, 457
  lanthanides from water with dibutylphenacylphosphonate, 257
  nickel(II) camphorquinone dioxime, 503
  nickel(II) β-diketonate complexes, 461
  osmium(VI),(VIII) between carbon tetrachloride and water, 175
  synergic effects, 387, 457, 461, 683
  uranium(VI) with phosphate esters, 683
  zinc(II) β-diketonate, 387
Solvents, effect on solvent extraction of chromium(III) β-diketonates, 383
Solvolysis, chromium(III) thiocyanate complex, 191

Spectra, *see* circular dichroism, circularly polarised fluorescence, electronic, ESR, fluorescence, IR, mass, NMR, optical rotatory dispersion, photoacoustic, Raman
Spin-lattice relaxation, cobalamins and cobinamides, 321
Spreading pressure, dinonylnaphthalenesulphonic acid, 541
Stability constants, *see* formation constants
Stannylenes, 826
Static *trans* effect, tungsten(VI) imide complexes, 561
Styrene, hydroformylation, 592
Substitution reactions
  cobalt(III), 203
  platinum(II), 201
Sulphate complexes, manganese(II), 327
Sulphonic acid, oxime complex, critical adsorption concentration, critical micelle concentration and interfacial tension, 453
2-(4-Sulphophenyl)phosphonate, zirconium(IV), 61
3-Sulphopropylphosphonate, zirconium(IV), 61
5-Sulphosalicylate complexes, terbium(III), 693
Sulphur, pyramidal, inversion barriers, 289
Sulphur(VI)
  alkyl fluoride compounds, synthesis, IR, mass and Raman spectra and reactions, 723
  2-bromo-2,2-difluoroethyl-, 723
  1-chlorotetrafluoroethylpentafluoro-, 723
  1-deuterotetrafluoroethylpentafluoro-, 723
  1-deutero-2,2,2-trifluoroethylpentafluoro-, 723
  2,2-difluorovinylpentafluoro-, 723
  pentafluoro-2,2,2-trifluoroethyl-, 723
Sulphur dioxide, reaction with metallacycle, 747
Synergic effects, in solvent extraction, 387, 457, 461, 683

T

Tantalacyclopentanes, 763, 768
Tantalum(0)
  tris(2,2′-bipyridyl)-, synthesis and magnetic properties, 209
  tris(1,10-phenanthroline)-, synthesis and magnetic properties, 209
Tantalum(V), pentachloro-, 209
Tartaric acid, Pfeiffer effect with lanthanide complexes, 195
Tellurium(IV)
  diphenyl(2-methoxybenzoato)methyl-, 13
  diphenyl(methyl)-3-nitrobenzoato-, 13
  organo compounds, synthesis, electric conductance, IR, mass and NMR (^{1}H) spectra, 13
Telluronium salts, 13
Terbium(III)
  L-aspartatodi(5-sulphosalicylato)-, 693
  chiral complexes, fluorescence and circularly polarised fluorescence spectra, Pfeiffer effect and formation constant, 195
  complexes with 2,2′-diaminoether,*N*,*N*,*N*′,*N*′-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N*′,*N*′- tetraacetate, 277
  dihydroxofluoro-, 239
  di(5-sulphosalicylato)-L-lactato-, 693
  di(5-sulphosalicylato)-L-malato-, 693
  di(5-sulphosalicylato)-L-mandelato-, 693
  fluorohydroxo-, 239
  hydroxycarboxylate salicylate complexes, formation constants, fluorescence and circularly polarised fluorescence spectra and luminescence dissymmetry factors, 693
  tris(dibutylphenacylphosphonato)-, 257
  tris(pyridine-2,6-dicarboxylato)-, 195
  tris(pyridine-2,6-dicarboxylato)-, dibenzoyl tartaric acid complex, 195
  tris(pyridine-2,6-dicarboxylato)-, diethyl tartrate complex, 195
  tris(pyridine-2,6-dicarboxylato)-, dimethyl tartrate complex, 195
  tris(pyridine-2,6-dicarboxylato)-, di(4-toluyl) tartaric acid complex, 195
  tris(pyridine-2,6-dicarboxylato)-, tartaric acid complex, 195
2,2′:6′,2′′-Terpyridyl, cobalt(II) and (III) complexes, 21
2,3,4,5-Tetrachlorophenyl, mercury(II) compounds, 243
2,3,4,6-Tetrachlorophenyl, mercury(II) compounds, 243

- 2,3,5,6-Tetrachlorophenyl, mercury(II) compounds, 243
- Tetracyanoethene, reaction with iron pentacarbonyl, 799
- *closo*-Tetra(η^5-cyclopentadienylcobalt)tetraborane(4), 133
- *closo*-Tetra(η^5-cyclopentadienylnickel)tetraborane(4), 133
- Tetrafluoroacetaldehyde, reaction with tetrafluoromethylamide ion, 129
- Tetrafluoroboric acid, 441
- Tetrafluoroethene, reaction with rhodium(I) complex, 581
- Tetrafluoroethyl compounds
  - rhodium(I), 574
  - sulphur(VI), 723
- Tetrafluoromethylamide ion, reactive intermediate, 129
- Tetrahydroborate, 570
- Tetrahydrofuran complexes
  - neodymium(III), 423
  - rhenium(IV), 209
  - samarium(III), 423
  - uranium(III), 417
- 3,3′,4,4′-Tetrahydroxydiphenylsulphone dianion complexes, aluminium(III), 313
- 1-Tetralone oxime complex, rhodium(I), 501
- Tetramethylammonium salts, polyphosphates, 69
- Tetraphenylbuta-1,3-diene compounds, indium(III), 549
- Tetraphenylphosphonium hydrogen dinitrate, 525
- Tetraphenylporphyrinate complexes, iron(III), 737
- Tetrapolyphosphate, magnesium and tetramethylammonium salts, 69
- Thallium(I)
  - cyclopentadienide, 423
  - undecahydrodicarba-*nido*-undecaborate(2)-, 77
- Theophyllinate complexes, copper(II), 143
- Thermogravimetric analysis
  - cadmium(II) arsine carboxylate complexes, 667
  - caesium trihalides, 215
  - cobalt(III) binuclear complexes, 21
  - copper(II) carboxylate complex, 679
  - copper(II) morpholine nitrate complexes, 437
  - magnesium azide hydrazine complex, 421
  - molybdenum(VI) and tungsten (VI) oxides, reactions with sodium chloride, 471
  - thiazolylthiourea complexes, 557
  - zinc(II) arsine carboxylate complexes, 667
- Thiazole complexes
  - antimony(III) and bismuth(III), 655
  - synthesis, electric conductance, IR spectra and thermogravimetric analysis, 557
- 1-[2′-Thienyl]-4,4,4-trifluoro-1,3-butanedionate complexes
  - gallium(III) and indium(III), 9
  - zinc(II), 387
- Thioacetamide complex, iron(II), 429
- Thiocyanate, reaction with platinum(II) complex, 201
- Thiocyanate complexes
  - chromium(II), 135
  - chromium(III), 191
  - cobalt(II), 377, 711
  - copper(II), 377
  - manganese(II), 327
  - mercury(II), 711
  - nickel(II), 377, 711
- Thiodiethyleneglycol-1,4-bis(dibutylphosphate) complex, uranium(VI), 683
- Thioether complexes, palladium(II) and platinum(II), 289
- η^2-Thioformyl complex, rhodium(III), 576
- Thiolate complexes, cadmium(II), mercury(II) and zinc(II), 157
- Thionitrosyl complexes, rhodium(I) and (III), 576
- Thionyl chloride, 479
- Thiosulphate, reaction with platinum(II) complex, 201
- Thiourea, reaction with platinum(II) complex, 201
- Thiourea complexes
  - iron(II), 429
  - synthesis, electric conductance, IR spectra and thermogravimetric analysis, 557
- Thorium(IV)
  - aminopolycarboxylate complexes, formation constants and potentiometric titrations, 827

  bis[*N*-(2′-carboxyphenyl)iminodiacetato]-, 827
  bisiminodiacetato-, 827
  N-(2′-carboxyphenyl)iminodiacetato-, 827
  [(ethylenedioxy)diethylenedinitrilo]tetraacetato-, 827
  iminodiacetato-, 827
Threonine complexes, copper(II), 371
Thulium(III)
  complexes with 2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate , 277
  toluene-4-sulphonate, 253
  tris(dibutylphenacylphosphonato)-, 257
Tin, organo compounds, book review, 845
Tin(II)
  biscyclopentadienedithiocarboxylato-, tetraethylammonium salt, 187
  bis(trifluoromethyl)-, 826
  cyclopentadienedithiocarboxylate complexes, synthesis, Moessbauer and NMR (^{1}H) spectra, 187
  dithiocarboxylate complex, 187
  iodide, 826
  iodo(trifluoromethyl)-, 826
  organo compounds, formation, Moessbauer and NMR (^{19}F) spectra, 826
Tin(IV)
  N-arylsalicylideneimine complexes, NMR (^{1}H, ^{13}C) spectra, 213
  bromotrimethyl-, 397
  t-butylmethyl(phenyl)triphenylmethyl-, 1
  chlorotrimethyl-, 397
  cyclopentadienedithiocarboxylate complexes, synthesis, Moessbauer and NMR (^{1}H) spectra, 187
  dichlorodimethylbis*N*-salicylideneaniline-, 213
  dichlorodimethylbis*N*-salicylideneanisidine-, 213
  dithiocarboxylate complex, 187
  heteropolytungstates, 283
  methyl(2-methyl-2-phenylpropyl)(phenyl)triphenylmethyl-, 1
  organo compounds, synthesis, chromatographic resolution, electronic, mass and NMR (^{1}H) spectra, 1
  organo halides, isotopic substitution, IR and Raman spectra, normal coordinate analysis, force constants and bond lengths, 397
  triscyclopentadienedithiocarboxylato-, tetraethylammonium salt, 187
Titanacyclobutane, 762
Titanacyclobutene, 761
Titanacyclopentadienes, 760
Titanacyclopentanes, 745, 749
Titanacyclopentenes, 744, 745
Titanium(0), tris(1,10-phenanthroline)-, 209
Titanium(II)
  aluminium(III) chloride complex, 393
  η^{6}-benzenebistetrachloroaluminate, 393
  chloride complexes, IR spectra, polymorphism and neutron diffraction structure, 393
  tetrachloroaluminate, 393
Titanium(IV), tetrachloro-, 209
TLC, cobalamin and cobinamide isomers, 321
Toluene-4-sulphonate, NMR (^{1}H) spectra and lanthanide coplexes, 253
4-Tolyl compounds, mercury(II), 711
Total luminescence spectra, *see* fluorescence spectra
Transfer hydrogenation, 584
Trans influence, *see* static trans effect
1,4,7-Triazaheptane, reaction with coordinated di-2-pyridyl ketone, 21
Tribromide, caesium salt, 215
8,9,12-Tribromooctahydrodicarba-*nido*-undecaborate(2) complexes, cobalt(III), 511
Tri-*n*-butylborate, 701
Tri-*n*-butylboron, 701
Tri-*n*-butylphosphine complexes
  cobalt(-I), 45
  molybdenum(0), 130
Tri-*t*-butylphosphine complex, molybdenum(0), 130
8,9,12-Trichlorooctahydrodicarba-*nido*-undecaborate(2) complexes, cobalt(III), 511

2,3,4-Trichlorophenyl, mercury(II) compounds, 243
2,4,6-Trichlorophenyl, mercury(II) compounds, 243
Tricyclo[5.2.10]deca-3,8-diene, hydroformylation, 593
Tricyclohexylphosphine complex, molybdenum(0), 130
Triethylammonium chloride, 535
Triethylarsine complexes, cobalt(-I) and iron(-II), 45
Triethylborate, 701
Triethylboron, 701
Triethyleneglycol, solvent extraction, 383
Triethyleneglycol-1,6-bis(dibutylphosphate) complexes, uranium(VI), 683
1,1,1-Triethylhydrazinium chloride, 535
Triethylorthoformate, dehydrating agent, 409
Triethylphosphine complexes
  cobalt(-I) 45
  gold(I), 105
  iron(-II), 45
  platinum(II), 89
Triethylsilane, 601
Trifluoroacetate complex, rhodium(I), 574
Trifluoroacetylacetonate, *see* 1,1,1-trifluoro-2,4-pentanedionate
2,2,2-Trifluoroethyl compound, sulphur(VI), 723
Trifluoromethanimine, reactions with alkali fluorides and tetrafluoroacetaldehyde, 129
Trifluoromethyl compounds, cadmium(II), tin(II), 826
Trifluoromethylsulphonate complex, rhodium(I), 573
1,1,1-Trifluoro-2,4-pentanedionate complexes
  chromium(III), 383
  gallium(III) and indium(III), 9
Tri-4-fluorophenylphosphine complexes, rhodium(I), 339
Triiodide, caesium salt, 215
8,9,12-Triiodooctahydrodicarba-*nido*-undecaborate(2) complexes, cobalt(III), 511
Trimetaphosphate
  reaction with ammonia, HPLC, kinetic study and NMR (^{31}P) spectra, 425
  sodium salt, trihydrate, 425
α,α′,α″-Trimethoxymesitylene, 563
Trimethoxymethylmesitylene, 563
Trimethoxymethylpseudocumene, 563
α,α′,α″-Trimethoxypseudocumene, 563
Trimethylborate, 701
Trimethylboron, 701
1,3-Trimethylenebis(diphenylphosphine selenide) complex, zinc(II), 781
1,3-Trimethylenebis(diphenylphosphine sulphide) complex, zinc(II), 781
Trimethylphosphine complexes
  osmium(II), (IV) and (II,III), 83
  rhenium(I), 31, 37, 441, 803
  rhenium(III), 37, 441
  rhenium(IV), 37
  rhenium(V), 31, 37, 441
  rhodium(I), 809
Trimethylphosphine oxide complexes, europium(III), gadolinium(III), indium(III), lanthanum(III), and praseodymium(III), 307
Trimethylphosphite
  complexes, copper(II), 467
  oxidation by copper(II), 467
Trimethylphosphonium, hexachloroosmate(IV) salt, 83
Trimethylsilylmethyl complexes
  osmium(III), (IV) and (VI), 83
  rhodium(I), 809
Trinuclear complex, rhenium(V,I,V), 31
Trioctylamine
  solvent extraction of metal chloro complexes, 623
  synergic effects in solvent extraction, 461
Trioctylammonium chloride, 457
Trioctylmethylammonium chloride, 457
Triphenylarsine complexes
  molybdenum(0), 130
  rhodium(I), 339
Triphenylboron, 701

Triphenylmethyl
  tetrafluoroborate, 441
  tin compounds, 1
Triphenylmethyl cation, reaction with metallacycle, 747
Triphenylphosphine, 535
  displacement from rhodium(I) complex, 569
  promoter of selective hydroformylation, 591
  solvent for hydroformylation, 595
Triphenylphosphine complexes
  gold(I), 105
  molybdenum(0), 130
  osmium(II), (III) and (IV), 83
  palladium(II), 539
  rhenium(V), 37
  rhodium(I), 339, 569, 572, 809
Triphenylphosphine selenide complexes, mercury(II), zinc(II), 781
Triphenylphosphine sulphide complexes, mercury(II), zinc(II), 781
Triphenylphosphite complexes
  cobalt(-I) and iron(-II), 45
  molybdenum(0), 130
  rhodium(I), 572
Triphenylphosphonium, hexachloroosmate(IV) salt, 83
Triphenylphosphoniumacetylmethylide complex, palladium(II), 539
Triphenylphosphoniumbenzoylmethylide complex, palladium(II), 539
Triphenylstibine complexes, molybdenum(0), 130
Triphenyltelluronium fluoride, 13
Tripolyphosphate, magnesium and tetramethylammonium salts, 69
Tri-*i*-propylboron, 701
Tri-*n*-propylborate, 701
Tri-*n*-propylboron, 701
Tri-*i*-propylphosphine complexes, rhodium(I), 839
Tri-*n*-propylphosphine complexes, cobalt(-I) and iron(-II), 45
Trisdimethylaminoboron, 701
Tris(diphenylthiophosphinyl)methanido complexes, cadmium(II) and mercury(II), 137
Tris(2-pyridyl)phosphine complex, rhodium(I), 572
Tri-3-tolylphosphine complexes, rhodium(I), 339
Tri-4-tolylphosphine complexes, rhodium(I), 339
Tri-4-tolylphosphine selenide complexes, cadmium(II), mercury(II), zinc(II), 781
Tri-4-tolylphosphine sulphide complexes, cadmium(II), zinc(II), 781
Trityl, *see* triphenylmethyl
Tungstacyclobutanes, 770
Tungstacyclopentanes, 750
Tungsten, removal from molybdenum, 479
Tungsten(0)
  bipyrimidine bridged binuclear complex, 53
  μ-2,2′-bipyrimidine(octacarbonyl)molybdenum-, 53
  carbonyl phosphine complexes, 819
  hexa-*t*-butylisocyanide-, 803
  isocyanide complex, synthesis, IR and NMR (^{1}H, ^{13}C) spectra, 803
  tris(2,2′-bipyridyl)-, 209
  tris(1,10-phenanthroline)-, 209
Tungsten(I)
  di(2-oxo-6-methylpyridinato)tetra-*t*-butylisocyanidedi-, 803
  isocyanide oxopyridinate complex, synthesis, IR and NMR (^{1}H, ^{13}C, ^{195}Pt) spectra, 803
Tungsten(II)
  cyclopentadienyl(methyl)tricarbonyl-, 663
  methyl(pentamethylcyclopentadienyl)tricarbonyl-, 663
  methyltricarbonyl[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
  tetra(2-oxo-6-methylpyridinato)di-, 803
Tungsten(IV)
  alkoxy alkyl cyclopentadienyl complexes, synthesis, IR, mass and NMR (^{1}H) spectra, 629
  benzoatobiscyclopentadienyl(methyl)-, 629
  (2,2′-bipyridyl)tetrachloro-, synthesis, 209
  biscyclopentadienylchloro(4-methylphenoxy)-, 629
  biscyclopentadienyldimethyl-, 629
  biscyclopentadienyl[dimethyl(phenyl)phosphine]hydrido-, hexafluorophosphate, 629
  biscyclopentadienylhydridomethyl-, 629
  biscyclopentadienyliodo(methyl)-, 629

  biscyclopentadienyl(methoxy)-4-methylphenyl-, 629
  biscyclopentadienyl-4-methoxyphenyl(methyl)-, 629
  biscyclopentadienyl(methyl)-4-methylphenoxy-, 629
  biscyclopentadienyl(methyl)phenoxy)-, 629
  biscyclopentadienyl(methyl)phenoxymethyl-, 629
  diacetonitriletetrachloro-, 209
  (1,10-phenanthroline)tetrachloro-, synthesis, 209
Tungsten(V)
  cyanide complex, IR and NMR (^{1}H) spectra, X-ray diffraction and phase transitions, 779
  hexaphenoxo-, bistetrahydrofuranlithium, potassium, sodium and tetraethylammonium salts, 641
  octacyano-, tetra-*n*-butylammonium salt, 779
  pentaphenoxo-, 641
  phenoxide complexes, synthesis, ESR, IR and NMR (^{1}H) spectra and X-ray crystal structure, 641
Tungsten(VI)
  alkoxide imide complexes, synthesis, NMR (^{1}H, ^{13}C) spectra, X-ray crystal structures and static *trans*- effect, 561
  t-butylaminechloro(phenylimido)tri-*t*-butoxo-, 561
  t-butylaminephenylimidotetra*neo*-pentoxo-, 561
  t-butylaminephenylimidotetra*i*-propoxo-, 561
  hexaphenoxo-, 641
  octamethoxotetraphenylimidodi-, 561
  phenylimidotetrachloro-, 561

U

Undecahydrodicarba-*nido*-undecaborate(2) complexes
  cobalt(III), 511
  comparison with pentamethylcyclopentadiene complexes, 663
Uranium(III)
  cyclopentadienyl compounds, synthesis, electronic, IR, mass and NMR (^{1}H) spectra and X-ray diffraction, 417
  μ-pyrazinedi(triscyclopentadienyl-), 417
  μ-pyrazinedi(trismethylcyclopentadienyl-), 417
  tetrahydrofuran(triscyclopentadienyl-), 417
Uranium(III) (cont)
  tetrahydrofuran(trismethylcyclopentadienyl)- , 417
Uranium(IV)
  bispentamethylcyclopentadienyldichloro-, 663
  bis[undecahydrodicarba-*nido*-undecaborate(2)]-, 663
  chloride, 417
  tetrakiscyclopentadienyl-, 663
Uranium(VI)
  alaninedioxo-, 537
  amminobis(1,1,1,5,5,5-hexafluoro-2,4-pentanedionato)dioxo-, 483
  ammonia diketonate oxo complex, colour change, X-ray crystal structure and hydrogen bonding, 483
  bisalaninedioxo-, 537
  bis[*N*-(2′-carboxyphenyl)iminodiacetato]dioxo-, 827
  biscysteinedioxo-, 537
  bis[{(ethylenedioxy)diethylenedinitrilo}tetraacetato]dioxo-, 827
  bis(iminodiacetato)dioxo-, 827
  bisisoleucinedioxo-, 537
  bisleucinedioxo-, 537
  bismethioninedioxo-, 537
  bisserinedioxo-, 537
  bis[thiodiethyleneglycol-1,4-bis(dibutylphosphate)]dinitratodioxo-, 683
  bisvalinedioxo-, 537
  N-(2′-carboxyphenyl)iminodiacetatodioxo-, 827
  cysteinedioxo-, 537
  dinitratodioxohemi[ethyleneglycol-1,2-bis(dibutylphosphate)]-, 683
  dinitratodioxohemi[diethyleneglycol-1,4-bis(dibutylphosphate)]-, 683
  dinitratodioxohemi[triethyleneglycol-1,6-bis(dibutylphosphate]-, 683
  dinitratodioxosesqui[ethyleneglycol-1,2-bis(dibutylphosphate)]-, 683
  dinitratodioxosesqui[diethyleneglycol-1,4-bis(dibutylphosphate)]-, 683
  dinitratodioxosesqui[triethyleneglycol-1,6-bis(dibutylphosphate)]-, 683

  dioxo-, aminoacid complexes, formation constants, 537
  dioxo-, aminopolycarboxylate complexes, formation constants and potentiometric titrations, 827
  dioxo-, apatites, synthesis, fluorescence, IR and Raman spectra, electron probe microanalysis, thermal decomposition and X-ray diffraction, 343
  dioxo[(ethylenedioxy)diethylenedinitrilo]tetraacetato-, 827
  dioxo-, fluoride complex, synthesis and electronic, fluorescence, IR and Raman spectra, 835
  dioxo(iminodiacetato)-, 827
  dioxoisoleucine-, 537
  dioxoleucine-, 537
  dioxo(methionine)-, 537
  dioxo-, nitrate phosphate ester complexes, solvent extraction, 683
  dioxo(serine)-, 537
  dioxotetrafluoro-, 835
Uranium(VI) (cont)
  dioxo(valine)-, 537
Uridine, reaction with chromium(VI), 497
5′-Uridinemonophosphate, reaction with chromium(VI), 497
UV spectra, see electronic spectra

V

Valine complexes, uranium(VI), 537
Vanadium, tris(2,2′-bipyridyl)nitrosyl-, synthesis and IR spectrum, 209
Vanadium(0)
  (2,2′-bipyridine)tetracarbonyl-, synthesis and IR spectrum, 209
  tris(2,2′-bipyridyl)-, 209
  tris(di-t-butyldiimine)-, synthesis and magnetic properties, 209
  tris(2,2′-dipyridylamine)-, synthesis and magnetic properties, 209
  tris(2,2′-dipyridyldisulphide)-, synthesis and magnetic properties, 209
  tris(1,10-phenanthroline)-, 209
Vanadium(II), bis(2,2′-bipyridyl)iodo(methyl)-, synthesis, 209
Vanadium(III) chloride, 209
Vanadium(IV), oxo-, sulphate, hydrated complex with poly(iminoethylene)dithiocarbamate, 225
Vanadium(V)
  diperoxofluoro(oxo)-, ammonium, caesium,potassium and rubidium salts, 553
  fluoro oxo peroxo complexes, synthesis and IR spectra, 553
  oxide, 553
Vinyl complexes, rhodium(I), 575
Vinyl cyanide, hydrosilylation, 600
Vinyldimethylcyclohexene, hydroformylation, 593
N-Vinylphthalimide, hydroformylation, 599
N-Vinylsuccinimide, hydroformylation, 599
Viscosity
  chloride complexes in nitrobenzene, 623
  osmium carbohydrate polymers, 335
Vitamin B_6 complexes, 791
Vitamin B_{12}, see cobalamins and cobinamides

W

Wade's rules, violation in dodecahedral cluster compounds, 133
Water, hydrolysis
  chromium(III) thiocyanate complex, 191
  platinum(II) halide complex, 201
Water, solvent extraction, 383, 387
Water complexes
  chromium(III), 471
  copper(II), 143, 247
  molybdenum(V), 109
  nickel(II), 409
  platinum(II), 57

X

X-ray crystal structure
  ammonium β-octamolybdate pentahydrate, 17
  caesium perbromate, 567

  chromium(III) chloride methylnicotinate water complex, 471
  cobalt(III) hexammine salt, 365
  copper(II) chloride complex, 365
  copper(II) nitroacetate complex, 247
  gold-gold bonded osmium cluster carbonyl hydride, 105
  hydrogen dinitrate salt, 525
  indium(III) chloride phosphine oxide complex, 355
  iron(0) cluster acyl carbonyl hydride complex, 842
  iron(0) cluster alkoxide alkylidyne carbonyl complex, 842
  iron(II) arene carbaborane complex, 77
  mercury(II) mercapto complexes, 647
  molybdenum(II) carbonyl cyclopentadienyl pyrrole Schiff base complex, 617
  nickel(II) chloride crown ether complex, 409
  osmium gold cluster carbonyl hydride, 105
  osmium(VI) acetate oxo complex, 840
  platinum(II) phosphine complex, 89
  rhenium(I) amido-complex, 37
  rhenium(I) chloride isocyanide phosphine complexes, 803
  rhenium(I) cyclopentadienyl phosphine complex, 441
  rhenium(I) dinitrogen hydride phosphine complex, 441
  rhenium(I) metallated phosphine complex, 441
  rhenium(III) chloride hydride phosphine complex, 441
  rhenium(III) iodide methyl phosphine complex, 441
  rhenium(V) amido-complex, 37
  rhodium(I) carbonyl hydride phosphine complex, 571
  rhodium(I) cyclopentadienyl phosphine complex, 809
  ruthenium(III) chloride pyridine complex, 525
  trinuclear rhenium(V,I,V) complex, 31
  tungsten(V) phenoxide complexes, 641
  tungsten(VI) alkoxide amine chloride imide complex, 561
  tungsten(VI) alkoxide imide complex, 561
  uranium(VI) ammonia diketonate oxo complex, 483
X-ray diffraction
  halide mercaptide complexes, cadmium(II), mercury(II) and zinc(II), 157
  heteropolytungstates, 283
  phenanthridine and phenanthridinium hydrogen dinitrate and nitrate, 525
  tungsten(V) cyanide complex, 779
  uranium apatites, 343
  uranium(III) cyclopentadienyl pyrazine complex, 417
  zirconium(IV) sulphophosphonates, 61

Y

Ylide complexes
  palladium(II), 539
Ylide complexes (cont)
  platinum(II) and (IV), 689
Ytterbium, reaction with thallium(I) cyclopentadienide, 423
Ytterbium(II)
  biscyclopentadienyl-, 423
  biscyclopentadienyl(1,2-dimethoxyethane)-, 423
  fluoride and oxide, improbability of disproportionation, 219
Ytterbium(3+), reaction with europium(III) complex, 707
Ytterbium(III)
  complexes with 2,2′-diaminodiethylether,*N*,*N*,*N*′,*N*′-tetraacetate and 1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate, 277
  difluorohydroxo-, 239
  dihydroxofluoro-, 239
  propylenediaminetetraacetato-, 707
  toluene-4-sulphonate, 253
  triscyclopentadienyl-, 423
Yttrium(III), complex with 1,5-diaminopentane,*N*,*N*,*N*′,*N*′-tetraacetate, 277

Z

Zerovalent compounds, convenient synthesis, 209
Zinc isotope (^{65}Zn) tracer study, 387
Zinc(II)
  alanine complex, 153

  aminothiazoline complex, 153
  arsine carboxylate complexes, synthesis, IR spectra and thermogravimetric analysis, 667
  biacetyldihydrazone complex, synthesis, IR spectrum and electric conductance, 831
  bis(*N*-benzothiazolyl-*N*′-ethylthiourea)-, 557
  bis(*N*-benzothiazolyl-*N*′-methylthiourea)-, 557
  bis(biacetyldihydrazone)dichloro-, 831
  bis(2-dicyclohexylarsinobenzoato)- trihydrate, 667
  bisdiethylamido-, 563
  bis(2-diethylarsinobenzoato)- monohydrate, 667
  bis-μ-(1-methyl-4-mercaptopiperidine)tetrabromodi-, monohydrate, 157
  bis-μ-(1-methyl-4-mercaptopiperidine)tetraiododi-, monohydrate, 157
  bis[1-(2′-thienyl)-4,4,4-trifluoro-1,3-butanedionato]-, 387
  bis[tetracarbonylcobaltate(-I)]-, 563
  bis(triphenylphosphine selenide)tetraiododi-, 781
  bis(triphenylphosphine sulphide)diiodo-, 781
  bis(tri-4-tolylphosphine selenide)tetraiododi-, 781
  bis(tri-4-tolylphosphine sulphide)tetraiododi-, 781
  chloride, 543
  cystinate complexes, 153
  diacetatodiaquo-, 557
  2,6-dibenzothiazol-2-ylpyridinedichloro-, 543
  dihistidine(pyridoxamine)-, 791
  diiodo[1,3-trimethylenebis(diphenylphosphine selenide)]-, 781
  diiodo[1,3-trimethylenebis(diphenylphosphine sulphide)]-, 781
Zinc(II) (cont)
  β-diketonate complex, solvent extraction in presence of heterocyclic diamines, 387
  halide phosphine selenide and phosphine sulphide complexes, synthesis, electric conductance and IR spectra, 781
  halide thiolate complexes, synthesis, IR spectra, electric conductance and X-ray diffraction, 157
  heteropolytungstates, 283
  histidine(pyridoxamine)-, 791
  histidine pyridoxamine complexes, formation constants and potentiometric titrations, 791
  iodide, 781
  manganese phosphate $[Mn_3Zn_9(PO_4)_8]$, 349
  tetrachloro-, methyltrioctylammonium and trioctylammonium salts, 623
Zirconacyclopentadiene, 744, 760
Zirconacyclopentanes, 749, 763, 768, 770
Zirconacyclopentenes, 745, 763
Zirconium(0)
  tris(2,2′-bipyridyl)-, 209
  tris(1,10-phenanthroline)-, 209
Zirconium(IV)
  bis(*N*,*N*′-disalicylidene-3,4-phenylenediamine-1-ethylbenzoato)-, methanol solvate, 487
  bis(η^5-pentamethylcyclopentadienyl)dihydrido-, 663
  bis[undecahydrodicarba-*nido*-undecaborate(2)]dihydrido-, 663
  (η^1-cyclopentadienyl)tri(η^5-cyclopentadienyl)-, 663
  di(η^5-cyclopentadienyl)dihydrido-, 663
  eight-coordinate tetradentate Schiff base complex, synthesis, electronic, IR, mass and NMR (^1H) spectra and ligand synthesis and isomerism, 487
  2-(4-sulphophenyl)ethylphosphonate, 61
  sulphophosphonates, synthesis, X-ray diffraction, DTA study, cation exchange reaction, catalytic properties, 61
  3-sulphopropylphosphonate, 61
  tetrachloro-, 209
  tetra-*n*-butoxo-, butanol solvate, 487

New and Forthcoming Reference Works from IUPAC

CHEMICAL DATA SERIES

Scientists, engineers and information managers will save hours of time searching the literature for physical and chemical data by first consulting the *Chemical Data Series.* Under the auspices of IUPAC, experts have collected data sought most frequently, such as thermodynamic and transport properties, equilibrium constants, stability constants and ionisation constants. Data in each volume are presented in tabular format with full references to the original work. In many cases the data have been critically evaluated by experts, paying particular attention to experimental conditions, techniques and estimated errors. Laboratories and reference libraries in the areas of chemistry, physics, materials science and engineering cannot afford to be without this highly acclaimed source of chemical data.

Recent and forthcoming volumes—

THERMODYNAMIC & TRANSPORT PROPERTIES OF ALKALI METALS

Edited by **R W OHSE & H BINDER,** *Joint Research Centre, Karlsruhe, Federal Republic of Germany*

X: *Chemical Data Series, Volume 30*

1500 pp approx (2 volumes) 1984

0 08 026176 0 (H)† **$350.00**

IONISATION CONSTANTS OF INORGANIC ACIDS & BASES IN AQUEOUS SOLUTION

2nd Edition

Edited by **D D PERRIN,** *Australian National University, Canberra*

X: *Chemical Data Series, Volume 29*

194 pp 1500 lit refs approx 1982

0 08 029214 3 (H) **$50.00**

STABILITY CONSTANTS OF METAL-ION COMPLEXES, Part A: INORGANIC LIGANDS

Edited by **E HÖGFELDT,** *Royal Institute of Technology, Stockholm, Sweden*

X: *Chemical Data Series, Volume 21*

324 pp 3000 lit refs approx 1982

0 08 020959 9 (H) **$85.00**

SOLUBILITY DATA SERIES

Editor-in-Chief: **A S KERTES,** *Hebrew University, Jerusalem, Israel*

The *Solubility Data Series* has become established as the most authoritative and comprehensive numerical work of reference ever published on the subject of solubility. When complete the Series will cover all solid, liquid and gaseous systems for which experimental data have been published.

Data are critically evaluated paying particular attention to experimental techniques and purity of reagents. This multidisciplinary reference set will be essential in all libraries which serve the information needs of researchers in chemistry, chemical engineering, the environmental sciences, biology, materials science, medicine, pharmacology and the geosciences.

Recent and forthcoming volumes –

SULFUR DIOXIDE, CHLORINE, FLUORINE & CHLORINE OXIDES

Edited by **C L YOUNG,** *University of Melbourne, Australia*

X: *Solubility Data Series, Volume 12*

496 pp 1500 lit refs approx Summer 1983

0 08 026218 X (H)† **$100.00**

NITROGEN & AIR

Edited by **R BATTINO,** *Wright State University, Dayton, OH, USA*

X: *Solubility Data Series, Volume 10*

618 pp 1500 lit refs approx 1982

0 08 023922 6 (H) **$100.00**

ETHANE

Edited by **W HAYDUK,** *University of Ottawa, Ontario, Canada*

X: *Solubility Data Series, Volume 9*

286 pp 750 lit refs approx 1982

0 08 026230 9 (H) **$100.00**

RECOMMENDED METHODS FOR PURIFICATION OF SOLVENTS & TESTS FOR IMPURITIES

Edited by **J F COETZEE,** *University of Pittsburgh, PA, USA*

This book brings together IUPAC's official recommendations for purifying the most important aprotic and amphiprotic solvents. It provides a critical evaluation of methods for purification and tests for impurities, and lists those physical and chemical properties that are particularly relevant in electrochemical applications. Information on the toxicity of solvents is also included. For analytical, organic and inorganic chemists, electrochemists and spectroscopists in universities, government and industrial laboratories.

Contents (partial): Solvent purity: general considerations. Relative reactivities of solutes in different solvents: guidelines to probable impurity effects. Acetonitrile. Sulfolane. Propylene carbonate. Dimethyl sulfoxide Dimethylformamide. Hexamethylphosphoramide Pyridine. Ethylenediamine. N-Methylacetamide N Methylpropionamide

X: *IUPAC Publication*

72 pp 2 illus 402 lit refs (R) 1982

0 08 022370 2 (F) **$20.00**

Prices are subject to change without notice
X: Books published in series. Standing/continuation orders are invited.

PERGAMON PRESS

Headington Hill Hall, Oxford OX3 0BW, UK
Fairview Park, Elmsford, New York, 10523, USA

Individual volumes now available

COMPREHENSIVE ORGANIC CHEMISTRY

The Synthesis and Reactions of Organic Compounds

Chairman of the Editorial Board: **Sir DEREK BARTON**, FRS, *CNRS, Gif-sur-Yvette, France*
Deputy Chairman: **W DAVID OLLIS**, FRS, *University of Sheffield, UK*

Independent Reviews from the World's Literature

"This is an important work, and one that chemical libraries cannot afford to be without." —**Journal of the American Chemical Society**

"It is indeed comprehensive and will be an essential purchase for chemical libraries." —**Nature**

"Undoubtedly every chemistry library should have a complete set." —**Chemistry in Britain**

"It should be in every chemistry library as a standard reference work." —**New Scientist**

"A highly valuable and well-presented source of authoritative and accurate information." —**Journal of Medicinal Chemistry**

"It is certain that this work will become indispensable as a source of reference." —**Synthesis**

"The extensive treatment of the newer methods of organic synthesis will be of great interest." —**Journal of Pharmaceutical Sciences**

"It will be the most useful single reference work for the next decade." —**Choice**

Volume 1 — STEREOCHEMISTRY, HYDROCARBONS, HALO COMPOUNDS, OXYGEN COMPOUNDS

Edited by **J F STODDART**, *University of Sheffield, UK*

Contents: Nomenclature and stereochemistry. Hydrocarbons. Halo compounds. Alcohols, phenols, ethers and related compounds. Aldehydes and ketones.

1242 pp (R) 1979
0 08 021313 8 (H) **$230.00**

Volume 2 — NITROGEN COMPOUNDS, CARBOXYLIC ACIDS, PHOSPHORUS COMPOUNDS

Edited by **I O SUTHERLAND**, *University of Liverpool, UK*

Contents: Amines and related compounds. Nitro and nitroso compounds. Imines, nitrones, nitriles and isocyanides. Carboxylic acids and related compounds. Phosphorus compounds.

1344 pp (R) 1979
0 08 021314 6 (H) **$230.00**

Volume 3 — SULPHUR, SELENIUM, SILICON, BORON, ORGANOMETALLIC COMPOUNDS

Edited by **D N JONES**, *University of Sheffield, UK*

Contents: Organic sulphur compounds. Organic selenium and tellurium compounds. Organic silicon compounds. Organic boron compounds. Organometallic compounds.

1338 pp (R) 1979
0 08 021315 4 (H) **$230.00**

Volume 4 — HETEROCYCLIC COMPOUNDS

Edited by **P G SAMMES**, *University of Leeds, UK*

Contents: Nitrogen systems — the azines. Nitrogen systems — the azoles and other systems. Oxygen systems. Sulphur and other heteroatom systems. Mixed heteroatom systems.

1242 pp (R) 1979
0 08 021316 2 (H) **$230.00**

Volume 5 — BIOLOGICAL COMPOUNDS

Edited by **E HASLAM**, *University of Sheffield, UK*

Contents: Biological chemistry — introduction. Nucleic acids. Proteins — amino acids and peptides. Proteins — enzyme catalysis and functional proteins. Lipid chemistry and biochemistry. Carbohydrate chemistry. Synthesis of organic macromolecules and their uses in organic chemistry. Bio-organic chemistry. Biosynthetic pathways from acetate. Biosynthesis — a general survey.

1220 pp (R) 1979
0 08 021317 0 (H) **$230.00**

Volume 6 — FORMULA, SUBJECT, AUTHOR, REACTION, & REAGENT INDEXES

Edited by **C J DRAYTON**, *Pergamon Press, Oxford, UK*

Formula index. Subject index. Author index. Reaction index. Reagent index.

1642 pp (R) 1979
0 08 022931 X (H) **£115.00 $230.00 US$264.50**

6 Volume Set

8028 pp (R) 1979
0 08 021319 7 (H) **$1375.00**

PERGAMON PRESS
Headington Hill Hall, Oxford OX3 0BW, UK
Fairview Park, Elmsford, New York 10523, USA

Pergamon launches new series of monographs

ORGANIC CHEMISTRY SERIES

Series Editor: J E BALDWIN, FRS, *University of Oxford, UK*

Pergamon is pleased to announce publication of the first volume in the new *Organic Chemistry Series* of monographs. The Series will endeavour to encompass modern and important themes in the field of organic chemsitry. Each volume will provide an approach to problem solving in synthesis and encourage a philosophical view of modern organic chemistry.

SYNTHETIC DESIGN WITH CHIRAL TEMPLATES

S HANESSIAN, *University of Montreal, Canada*

This book explains and illustrates a new strategy for the synthesis of natural products using optically active starting materials in general and carbohydrates in particular. The strategy involves recognizing symmetry elements, carbon framework patterns and functional group interrelations in the desired product and locating a suitable 'chiral template' derived from a carbohydrate. A short introduction covers the philosophy of using carbohydrates for enantiospecific synthesis and enumerates the ground rules and principles involved. The greater part of the book reviews the many syntheses of natural products starting from simple sugars. Excellent use is made of flow sheets. Carbohydrates emerge as surprisingly versatile 'building blocks' for the molecular architect. Of interest to researchers and students in organic chemistry.

Contents: Basic Concepts: Stereochemistry, stereoelectronic effects and conformation. Symmetry relationships. Design: The concept of 'chiral templates'. Types of targets derived from carbohydrate precursors by synthesis. Discovery: Where in the carbon skeleton is the carbohydrate? Execution: Molecules containing apparent carbohydrate-type symmetry. Molecules containing partially hidden carbohydrate-type symmetry. Molecules containing hidden carbohydrate-type symmetry. References.

X: *Organic Chemistry Series, Volume 3*

200 pp approx 400 lit refs approx (R) Autumn 1983
0 08 029247 X (H)† **$40.00**

Just Published

ORGANOTRANSITION METAL CHEMISTRY: APPLICATIONS TO ORGANIC SYNTHESIS

S G DAVIES, *University of Oxford, UK*

This unique book provides an introduction to organometallic chemistry for chemists who have little or no experience in the field, describes the work that has already been done using organotransition metal complexes for synthesis, and examines the types of synthetic problems that can be solved using organometallic compounds. It covers all major advances that have been made in this rapidly expanding area of chemistry, including not only reactions with well established applications for synthesis but also many recently discovered potentially useful reactions, surveying recent literature. For advanced university students, organometallic chemists and organic chemists.

Contents: General introduction to organometallic chemistry. Complexation and decomplexation reactions. Organometallics as protecting and stabilising groups. Organometallics as electrophiles. Organometallics as nucleophiles. Coupling and cyclisation reactions. Isomerisation reactions. Oxidation and reduction. Carbonylation and related reactions. Index.

X: *Organic Chemistry Series, Volume 2*

428 pp 600 illus approx 648 lit refs (R) 1982
0 08 026202 3 (H) **$85.00**

STEREOELECTRONIC EFFECTS IN ORGANIC CHEMISTRY

P DESLONGCHAMPS, *University of Sherbrooke, Quebec, Canada*

Experimental evidence indicates that the geometry and reactivity of many organic molecules are controlled by stereoelectronic effects. These electronic effects are caused by the non-bonded electron pairs on heteroatoms such as oxygen, nitrogen and sulfur, and in particular by their orientation in space. This book introduces these new concepts, describes the effects on reactivity and presents the experimental evidence. Indispensable to organic chemists and biochemists in universities and industry and students preparing to enter these fields.

Contents: Introduction. Acetals and related functions. Esters and related functions. Amides and related functions. Reactions at saturated carbons. Reactions on SP_2 type unsaturated systems: nucleophilic addition; double-bond formation. Reactions on triple-bonds. Other reactions. Biological processes. Author index. Subject index.

X: *Organic Chemistry Series, Volume 1*

365 pp approx (R) May 1983
0 08 026184 1 (H) **$55.00**
0 08 029248 8 (F) **$27.50**

Prices are subject to change without notice.
X: Books published in series. Standing/continuation orders are invited.

PERGAMON PRESS

Headington Hill Hall, Oxford OX3 0BW, UK
Fairview Park, Elmsford, New York 10523, USA

Instructions to Contributors

1. CONTENT

Polyhedron will publish original fundamental experimental and theoretical work of the highest quality in the major areas of inorganic and organometallic chemistry. These areas include synthetic and reaction chemistry, kinetics and mechanisms of reactions, bioinorganic chemistry, photochemistry, and the use of metal and organometallic compounds in stoichiometric and catalytic synthesis of organic compounds.

Contributions will be rigorously refereed. They are to comprise full Papers, Notes and Preliminary Communications to the Editor. Publications will be in English. In addition, specially commissioned review articles (**Polyhedron Reports**) and collections of original research papers (**Polyhedron Symposia-in-Print**) of appropriate timeliness will be published. Every effort will be made to expedite publication and thus minimize the time delay between submission of an acceptable manuscript and its appearance in print.

2. PAPERS

These are to be substantial completed pieces of original research in inorganic and organometallic chemistry which have special significance, general interest and which, in addition, are presented clearly and concisely. An abstract is to be provided. Polyhedron papers will be printed either by conventional typesetting or by a photo-offset process (see below). The latter alternative will allow the publication of specially prepared manuscripts within 3 months of their acceptance. There has been an increasing tendency for important advances in inorganic and organometallic chemistry to be reported in the form of preliminary communications, in which space restrictions do not allow experimental details to be included; frequently these experimental details are published years later, if at all. Polyhedron now offers authors a medium for a rapid publication without rigid space limitations or any page charge.

3. NOTES

These are smaller reports on a more limited subject matter than papers but must be complete in themselves. A brief abstract should be provided. Polyhedron notes will mainly be printed by a photo-offset process (see Section 8), to achieve rapid publication.

4. COMMUNICATIONS

These are to be contributions of a preliminary nature and to be of length not normally exceeding 1000 words. They must contain novel and original findings which are of general interest and which will be of use to other workers in their field. This section will also provide a form for advancing or exchanging significant new interpretations of published data for which normal papers or notes would be inappropriate. Rapid publication will be achieved by using a photo-offset printing process (see Section 8).

5. POLYHEDRON REPORTS

These will be specially commissioned critical reports reviewing research results of topical importance. Such articles will be commissioned by the Editorial Board from recognized authorities in the appropriate fields.

6. POLYHEDRON SYMPOSIA-IN-PRINT

These will comprise collections of original research papers (including experimental sections) covering timely areas of inorganic and organometallic chemistry. Topics for forthcoming Symposia are announced from time to time. For each, a Symposium Editor will invite authors active in the field to submit papers. These will be rapidly reviewed and processed for publication by the Symposium Editor under the usual refereeing system.

Authors not already invited who have recent, significant results in the area announced may also submit contributions within a six-month deadline for Editorial consideration and possible inclusion.

7. PRESENTATION OF MANUSCRIPTS FOR TYPESETTING

Authors are requested to submit two copies of each manuscript. After the title of the paper and the authors' names prefixed by initials the following should be given: (i) establishment where the work was done; (ii) a space for insertion of the date the paper was received; (iii) an abstract of the paper. The authors to whom correspondence should be addressed by readers of the journal should be indicated by an asterisk.

The presentation of tables, graphs and formulae should be kept to an absolute minimum and where possible the results from a series of investigations should be in table form or as a graph. The same data should not be presented in both tables and figures or given in the Introduction as well as the Experimental Section.

8. PREPARATION OF TYPESCRIPTS FOR PHOTO-OFFSET PROCESS

The typescript, which will be photo-reduced by 25%, should be typed (preferably on an electric machine with 12 pitch type and a black carbon ribbon) on one side of the paper only within an area 19.7 cm wide by 31 cm deep and with margins of at least 4 cm. The typing should be in a single column, $1\frac{1}{2}$ spaced between lines. Any good quality paper may be used, but special size paper is available from any Pergamon office. Manuscripts for photo-offset printing must consist of an even number of pages; manuscripts of 3, 5, 7, etc. pages will be returned to the authors for shortening. Authors should also refer to Section 7 when preparing their typescript.

9. X-RAY CRYSTALLOGRAPHIC WORK

Manuscripts reporting X-ray crystallographic work must be accompanied by a list of refined coordinates for deposition at the Cambridge Crystallographic Data Centre (see detailed Announcement to Authors on p. ii of this issue).

10. TABLES

In the interests of economy and in order to avoid the introduction of errors, all tables will be reproduced by photo-offset means directly from the author's typed manuscript. This will materially reduce costs of publication and reduce setting delays. Tables should be submitted in a form suitable for direct reproduction. The following guidelines should be followed: (i) maximum page size should be 19.7 × 31 cm; (ii) large or long tables should be typed on continuing sheets; (iii) table headings should be typed on separate sheets but identifying numbers should be placed on the upper right-hand corner of each sheet of tabular material; (iv) **original typed tables are required**. It will be the author's responsibility to prepare his tables in accordance with this procedure as no facilities exist at the Editorial Office for retyping.

11. NOMENCLATURE

All nomenclature should be consistent, clear and unambiguous and should conform as closely as possible to the rules established by IUPAC. The following IUPAC publications (both available from Pergamon Press) should be consulted: "*Nomenclature of Inorganic Chemistry*, 1970 edition" and "*Nomenclature of Organic Chemistry*, 1979 edition".

12. FIGURES, FORMULAE, SCHEMES, ETC.

These should all be carefully prepared line drawings on plain white drawing paper, suitable for direct photographic reproduction. Originals are preferable, but photographs are acceptable. Photocopies are not acceptable. Each illustration should be clearly marked on the back with the names of the authors. Where necessary, the top of the figure should be indicated. Figure captions should be typed on a separate sheet.

13. REFERENCES

Text references to the literature should be by superscript numbers. The references themselves should be collected in order at the end of the manuscript. Journal abbreviations should conform to the "Chemical Abstracts Service Source Index".

14. MAILING OF MANUSCRIPTS

Papers, Notes and Communications

All manuscripts should be sent (in duplicate) to:

For U.S.A. and Canada

Professor M. H. Chisholm, Department of Chemistry, Indiana University, Bloomington, IN 47405, U.S.A.

For Rest of World

Professor D. C. Bradley FRS, Department of Chemistry, Queen Mary College, University of London, Mile End Road, London E1 4NS, UK

Polyhedron Symposia-in-Print

To the Symposium Editor as announced for the particular symposium.

Polyhedron Reports

Prof. M. H. Chisholm (for North, Central and South America), Department of Chemistry, Indiana University, Bloomington, IN 47405, USA

Dr. B. F. G. Johnson (for the Rest of the World), University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW, UK

15. REFEREEING OF PAPERS

All papers submitted are refereed by experts who advise the Executive Editor on the matter of acceptance in accordance with the high standards required and on the understanding that the subject matter has not been previously published and that the authors accept full responsibility for the factual accuracy of the data presented and have obtained any necessary authority to publish. The referees' names are not disclosed but their views are forwarded by the Executive Editor to the authors for consideration. All papers if accepted are subject to editorial amendment if necessary.

16. PROOFS

For typeset Papers, proofs together with reprint order form are posted by air, direct from the Printers to the chief author (indicated on the paper when submitted) for his approval and correction. These corrected proofs should be returned airmail without delay to Professor Bradley for inclusion in the next possible issue.

Authors are kindly requested when proof reading to correct only those errors introduced by the printer. Modification to the text at proof stage automatically causes delay in publication and incurs additional costs which may have to be charged to the authors.

17. REPRINTS

For typeset Papers, a reprint order form will be sent with the proofs. Authors submitting manuscripts for direct reproduction should indicate the number of reprints required when sending the manuscript otherwise only 25 free reprints will be sent.

18. BOOK REVIEWS, CONFERENCE ANNOUNCEMENTS

Appropriate books for review and full details of forthcoming meetings should be sent to the Executive Editor for possible inclusion.

AVAILABLE FOR IMMEDIATE DELIVERY

Comprehensive Organometallic Chemistry

The Synthesis, Reactions and Structures of Organometallic Compounds

Editor: Sir GEOFFREY WILKINSON, FRS Deputy Editor: F G A STONE, FRS Executive Editor: E W ABEL

Quotations from the reviews published in Tetrahedron Letters, 1982, Vol 23, No 44:

Supplied complete in nine beautifully bound volumes. Approximately 9500 pages, size 270x188mm.

The most authoritative and comprehensive work of reference in modern organometallic chemistry.

Written by a team of over 100 internationally recognised contributors.

Over 40,000 carefully selected literature references, providing the reader with a rapid directive towards the key sources in the organometallic chemistry literature.

Volumes 1 to 6 represent the first complete treatment of the organic chemistry of the metals, classified according to the Group of the Periodic Table. Here the editors have taken great care to identify significant trends and have allocated space accordingly. Important general topics such as Bonding Trends and Structural Correlations among Main Group Organometallic Compounds, Non-rigidity in Organometallic Compounds, Environmental Aspects of Organometallic Chemistry, and Bonding of Unsaturated Organic Molecules to Transition Metals are surveyed in separate sections.

Volumes 7 and 8 are devoted exclusively to the use of organometallic compounds in organic synthesis and in catalysis. In addition to accounts of the uses in organic synthesis of the organometallic compounds of a wide range of both main group and transition metals, there are specialist chapters on several topics.

Volume 9 is devoted exclusively to indexes, and covers in detail the whole subject matter of Volumes 1 to 8 with Subject, Formula and Author Indexes, along with two additional compilations: an Index of Organometallic Structures Determined by Diffraction Methods, and an Index of Review Articles and Specialist Texts, systematically arranged and covering the entire organometallic literature.

9-volume set Approximately 9500 pages
ISBN 0 08 025269 9 **US$2150.00 £1075.00**

Prices are subject to change without notice. Sterling price applies to customers in the UK and Eire only.

F A Cotton, *Texas A&M University, USA*
"It is a stupendous accomplishment on the part of editors, authors and publisher. In all its heroic sweep it never forgets about being concrete, useful and readable. It is, in short, magnificent. Perhaps libraries in big research institutions should purchase two sets, since heavy use may well lead to traffic problems in getting at only one."

E J Corey, *Harvard University, USA*
". . . will prove to be of unusual interest and value to students and practitioners of synthetic organic chemistry. A vast amount of useful information is here collected in beautifully organized and succinctly delivered form. The presentations are the handwork of leaders and experts in each field, as is evident from the great care which has been exercised in the selection of material and in the strong coverage of recent developments."

J F Normant, *Université Pierre et Marie Curie, France*
"Whatsoever from these various chapters I read, the leading conclusion that emerges is that the authors are excellent, and guarantee a high standard which will be most appreciated by chemists, academic as well as industrial, and I intend to let people know of it around me."

R Ugo, *University of Milan, Italy*
"Without any doubt this is the most authoritative and comprehensive work of reference on organometallic chemistry, which provides information to scientists, both in teaching and in industry, because great emphasis is given to industrial applications. It is expected that this series will become a highly commendable collection of material which will be an important reference work to chemists of all kinds."

H Nöth, *Universität München, Federal Republic of Germany*
"The editors are to be congratulated on choosing these authors, all of which presented excellent contributions."

G Wilke, *Max-Planck-Institut für Kohlenforschung, Federal Republic of Germany*
"It is my impression that the complete work will be excellent and extremely useful for all organometallic chemists. Already I would like to congratulate the publisher."

Free copies of the full text of these reviews, together with a brochure describing the work, will be supplied on request.

PERGAMON PRESS

Headington Hill Hall, Oxford OX3 0BW,
Fairview Park, Elmsford, New York 10523, USA

12/82/5D/1

Printed in Great Britain by A. Wheaton & Co. Ltd., Exeter